

4-1-2013

# Validating The Use Of Sex Steroid Hormones Extracted From The Skeletal Muscle Tissue Of The Atlantic Sharpnose Shark (*Rhizoprionodon Terraenovae*), Spiny Dogfish (*Squalus Acanthias*), And The Little Skate (*Leucoraja Erinacea*) To Assess Reproductive Status In Elasmobranch Fishes

Bianca Prohaska  
*University of New England*

Follow this and additional works at: <http://dune.une.edu/theses>

 Part of the [Marine Biology Commons](#)

© 2013 Bianca Prohaska

---

## Preferred Citation

Prohaska, Bianca, "Validating The Use Of Sex Steroid Hormones Extracted From The Skeletal Muscle Tissue Of The Atlantic Sharpnose Shark (*Rhizoprionodon Terraenovae*), Spiny Dogfish (*Squalus Acanthias*), And The Little Skate (*Leucoraja Erinacea*) To Assess Reproductive Status In Elasmobranch Fishes" (2013). *All Theses And Dissertations*. 11.  
<http://dune.une.edu/theses/11>

VALIDATING THE USE OF SEX STEROID HORMONES EXTRACTED FROM  
THE SKELETAL MUSCLE TISSUE OF THE ATLANTIC SHARPNOSE SHARK  
(*RHIZOPRIONODON TERRAENOVAE*), SPINY DOGFISH (*SQUALUS ACANTHIAS*),  
AND THE LITTLE SKATE (*LEUCORAJA ERINACEA*) TO ASSESS  
REPRODUCTIVE STATUS IN ELASMOBRANCH FISHES

BY

Bianca Karoly Prohaska B.S. Florida Institute of Technology, 2010

THESIS

Submitted to the University of New England in Partial Fulfillment of the Requirements  
for the Degree of

Master of Science


In

Marine Sciences

May, 2013


© 2013 Bianca Prohaska  
All Rights Reserved

This thesis has been examined and approved.




---

Thesis Director, Dr. James Sulikowski  
Associate Professor of Marine Science  
University of New England



---

Dr. A. Christine Brown  
Chair and Professor of Biology  
University of New England



---

Dr. Paul Tsang  
Professor of Molecular, Cellular And  
Biomedical Sciences  
University of New Hampshire

5/15/13

---

Date

## Acknowledgements

A sincere thank you to the many individuals that helped me throughout my Master's endeavors at the University of New England. First, I would like to acknowledge the captain's and crew of the fishing vessels that we collected our samples on, the *Lady Victoria*, *Mystique Lady*, *Proud Mary*, *Kimberly Grace* and *Elizabeth Helen*, as well as the captain and crew of the NOAA research vessel *Oregon II*. Thank you to all of the researchers at the NOAA Pascagoula Laboratory, especially Dr. William Driggers III, Dr. Eric Hoffmayer, Kristin Hannan, and Travis Holland. Additionally, thank you to all of the researchers at the Gulf Coast Research Lab, especially Jeremy Higgs, Jill Hendon, and Sarah Ashworth. Thank you so much to all of the members of the Sulikowski lab for their help in collecting and processing samples. I would also like to personally acknowledge Carolyn Wheeler for all of her time and effort in helping me finish up my lab work this last year at UNE. I especially want to thank my major professor, Dr. James Sulikowski, for his continual support and guidance throughout my thesis work, for giving me so many opportunities over and above my thesis, and for pushing me beyond my comfort zone, which allowed me to develop into the researcher that I am today. I also would like to thank my thesis committee members Dr. Paul C. W. Tsang, and Dr. A. Christine Brown for their supportive input. Paul, thank you so much for letting me spend time in your lab, it was truly amazing to learn radioimmunoassay from the "master". Thank you to Dr. Steve Travis for his availability to discuss the results of my statistical analyses. I greatly appreciate the support and encouragement of the Marine Science Center staff- Marian Reagan, Tim Arienti, Rick Miller, and Shaun Gill. Finally, I would

like to thank my family as well as my boyfriend Nick Noel IV for their support, encouragement and reassurance throughout this process. This work was sanctioned by the UNE's Animal Care and Use Committee (IACUC Protocol Number: UNE-20121107SULIJ). This thesis work was primarily supported by a Save Our Seas Foundation grant awarded to Dr. James Sulikowski, as well as graduate student funds from the Marine Science Department at UNE.

## TABLE OF CONTENTS

TITLE PAGE .....	i
COPYRIGHT .....	ii
THESIS APPROVAL .....	iii
ACKNOWLEDGEMENTS .....	iv
TABLE OF CONTENTS .....	vi
ABSTRACT .....	ix

### CHAPTER

### PAGE

I. CHAPTER 1: DEVELOPMENT OF A NON-LETHAL AND MINIMALLY		
INVASIVE PROTOCOL TO STUDY ELASMOBRANCH REPRODUCTION .....		11
Abstract .....		11
Introduction .....		12
Materials and Methods .....		14
Specimen Collection .....		14
Sampling .....		15
Plasma Steroid Hormone Extraction .....		16
Skeletal Muscle Tissue Hormone Extraction .....		16
Radioimmunoassay .....		18
Statistical Analysis .....		19
Results .....		19
<i>S. acanthias</i> .....		20

<i>R. terraenovae</i> .....	21
Discussion.....	22
<i>S. acanthias</i> .....	22
<i>R. terraenovae</i> .....	24
Conclusions.....	25
Acknowledgements.....	26
References.....	27
Tables.....	34
Table 1.....	34
Table 2.....	35
Figure Legends.....	36
Figure 1.....	37
Figure 2.....	38
Figure 3.....	39

II. CHAPTER 2: ASSESSING REPRODUCTIVE STATUS IN ELASMOBRANCH

FISHES USING STEROID HORMONES EXTRACTED FROM SKELETAL MUSCLE

TISSUE.....	40
Abstract.....	40
Introduction.....	41
Materials and Methods.....	44
Specimen Collection.....	44
Sampling.....	45
Plasma Steroid Hormone Extraction.....	45



Skeletal Muscle Tissue Hormone Extraction.....	46
Radioimmunoassay.....	47
Statistical Analysis.....	48
Results.....	49
<i>R. terraenovae</i> .....	49
<i>L. erinacea</i> .....	51
<i>S. acanthias</i> .....	53
Discussion.....	54
Conclusions.....	59
Acknowledgements.....	60
References.....	60
Tables.....	68
Table 1.....	68
Table 2.....	69
Table 3.....	70
Table 4.....	71
Table 5.....	72
Table 6.....	73
Figure Legends.....	74
Figure 1.....	76
Figure 2.....	77
Figure 3.....	78
III. APPENDIX.....	79

## ABSTRACT

VALIDATING THE USE OF SEX STEROID HORMONES EXTRACTED FROM THE SKELETAL MUSCLE TISSUE OF THE ATLANTIC SHARPNOSE SHARK (*RHIZOPRIONODON TERRAENOVAE*), SPINY DOGFISH (*SQUALUS ACANTHIAS*), AND THE LITTLE SKATE (*LEUCORAJA ERINACEA*) TO ASSESS REPRODUCTIVE STATUS IN ELASMOBRANCH FISHES

by

Bianca Prohaska

University of New England, May 2013

Currently, circulating concentrations of plasma steroid hormones are used as one means to assess reproductive maturity and reproductive cycles in elasmobranchs. However, obtaining blood non-lethally from large and/or endangered species can be problematic because of difficulties involved with specimen handling. Thus, there is a need to develop new approaches and techniques to study the reproductive biology of elasmobranchs. Previous work conducted on other classes of vertebrates has demonstrated that steroid hormones can be successfully extracted from muscle tissue. The process of collecting muscle tissue samples is quick, minimally invasive, and may be conducted without removing the animal from the water, facilitating its use on larger, and/or endangered species of elasmobranchs. Thus, the objective of the current study was to develop and validate a method for extracting steroid hormones from the skeletal muscle tissue of the lecithotrophic aplacental viviparous spiny dogfish (*Squalus acanthias*), the oviparous little skate (*Leucoraja erinacea*), and the placental viviparous Atlantic sharpnose shark (*Rhizoprionodon terraenovae*). The results suggest that

concentrations of progesterone [P<sub>4</sub>], testosterone [T], and estradiol [E<sub>2</sub>] present in skeletal muscle of the three aforementioned species can be successfully quantified by radioimmunoassay. Additionally, there were significant correlations between plasma and muscle E<sub>2</sub> concentrations in *S. acanthias*, as well as plasma and muscle P<sub>4</sub>, T and E<sub>2</sub> concentrations in *R. terraenovae* and *L. erinacea*. Furthermore, concentrations of muscle P<sub>4</sub>, T and E<sub>2</sub> were determined to be statistically significant indicators of reproductive status. The results of the current investigation demonstrated that skeletal muscle, which can be non-lethally harvested, is a viable and practical alternative to collecting blood samples for studying the reproductive biology of elasmobranchs.

## CHAPTER 1

### DEVELOPMENT OF A NON-LETHAL AND MINIMALLY INVASIVE PROTOCOL TO STUDY ELASMOBRANCH REPRODUCTION

#### Abstract

An understanding of basic reproductive biology is essential for successful species-specific management of elasmobranch fishes (sharks, skates, and rays). Such information is often gained through gross dissection or other lethal techniques, which are not appropriate for threatened and endangered species. Previous work on other vertebrates suggests that sex steroid hormones can be extracted from muscle tissues to identify reproductive status. The process of collecting a muscle biopsy is quick, minimally invasive, and can be collected without removing an animal from the water. Thus, the objective of the current study was to determine the efficacy of muscle steroid hormones to assess the reproductive biology of elasmobranch fishes. The results from the current study suggest that concentrations of muscle progesterone, testosterone and estradiol can be successfully quantified to study reproduction by radioimmunoassay. Additionally, there were significant correlations between plasma and muscle estradiol concentrations in *Squalus acanthias*, and progesterone, testosterone and estradiol concentrations in *Rhizoprionodon terraenovae*. The results of the present investigation demonstrated that skeletal muscle is a non-lethally harvested tissue that is well suited for studying the reproductive biology of elasmobranchs.

## **Introduction**

An essential component for assessing and managing populations of elasmobranchs (sharks, skates, and rays) is determining the species' specific reproductive biology, e.g. when sexual maturity occurs, timing of seasonal cycles, or gestation length (Walker 2004, 2005). Such information is essential for stock assessment purposes. Despite the importance of species-specific reproductive data to fisheries management (Walker 2004), these data are still unavailable for many species of elasmobranchs (Castro et al. 1999; Walker 2004; Pinhal et al. 2008), which has resulted in approximately 40% of the elasmobranch population being classified as data deficient (IUCN 2011). The lack of life history data when combined with direct and indirect fishing pressures make elasmobranchs susceptible to overexploitation worldwide, which can lead to population declines and localized extinction (Dulvy et al. 2003).

Historically, studies on the reproductive biology of elasmobranchs entailed sacrificing a relatively large number of specimens and observing changes in gross reproductive morphology over the reproductive cycle (e.g. Hisaw and Albert 1947; Zeiner and Wolf 1993; Walmsley-Hart et al. 1999; Francis et al. 2001; Ebert 2005; Ruocco et al. 2006). While this approach is still currently appropriate for elasmobranchs with sustainable population sizes (Heupel and Simpfendorfer 2010), it is not suitable for many because of their relatively low or unknown species abundance (IUCN 2011; Hammerschlag and Sulikowski 2011). Because of this limitation, alternative methods for determining size at maturity and/or reproductive cycles in elasmobranchs are being developed and validated (Sulikowski et al. 2007b).

One particular approach that has been utilized to non-lethally study the reproductive biology of elasmobranchs is the analysis of blood steroid hormone concentration (e.g. Perez and Callard 1993; Sulikowski et al. 2006; Kneebone et al. 2007; Sulikowski et al. 2007 a, b; Awruch et al. 2008; Henningsen et al. 2008). Reproductive hormones directly correlate with morphological changes within the reproductive tract, specifically at maturation (e.g. Gelsleichter et al. 2002; Sulikowski et al. 2006) and during reproductive cycles (e.g. Kneebone et al. 2007; Sulikowski et al. 2007 a, b). However, obtaining blood from large species is impractical and can be problematic because of difficulties involved with specimen handling. These restrictions also serve to protect the animals from stresses induced by handling procedures on large species (Skomal 2007). This is further compounded by the restrictions associated with sample collection from threatened and endangered species. Although plasma steroid hormone analysis is particularly useful for studying the reproductive biology of elasmobranchs, the aforementioned constraints make it necessary for new, less invasive approaches to be developed, especially for studying vulnerable species.

While steroid hormones are easily isolated from blood plasma, they are present in other body depots (Hoffmann 1978). For example, steroid hormones have been extracted from milk and/or muscle tissues from cows (Noppe et al. 2008), sheep, and chickens (Sawaya et al. 1998), urine and feces of free-ranging wildlife (Lasley and Kirkpatrick 1991; Holt and Pickard 1999), blubber from marine mammals (Mansour et al. 2002; Kellar et al. 2006) and the skeletal muscle tissue of chimaera (Barnett et al. 2009), for various reproductive analyses. Despite the successful extraction of steroid hormones from body depots other than plasma, this approach has never been conducted on an

elasmobranch. Given the lack of knowledge pertaining to the reproductive biology of elasmobranch fishes, the objectives of the present study were to determine: 1) if reproductive steroid hormones could be quantified from elasmobranch skeletal muscle tissue; and 2) if there is a relationship between muscle and plasma concentrations of these reproductive hormones.

## **Materials and Methods**

### **Specimen Collection**

Based on their relatively high abundance and ease of collection, the aplacental viviparous spiny dogfish (*Squalus acanthias*) and yolk sac placental viviparous Atlantic sharpnose sharks (*Rhizoprionodon terraenovae*) were selected as study organisms. Female *S. acanthias* were primarily captured by otter trawl off the coast of Rhode Island (USA) aboard the 13.7 m bottom trawler *F/V Proud Mary* between December 2010 and February 2012, in an area centered around 71.600 °W and 41.365 °N. Additional spiny dogfish samples were captured by gill net in the Gulf of Maine aboard the 14 m bottom trawler *F/V Lady Victoria* in November 2010, in an area centered around 70.257 °W and 42.917 °N. Female *R. terraenovae* were captured by bottom longline in the Gulf of Mexico aboard the 51.8 m *R/V Oregon II*, during the 2011 NOAA Bottom Longline Survey, in an area centered around 82.628 °W and 25.822 °N (Figure 1).

## **Sampling**

Immediately after capture, an 8 mL aliquot of blood was collected by caudal venipuncture using a heparinized needle and vacutainer. Blood samples were placed in a cooler (4°C) for up to 24 h before hematocrit analysis and centrifugation at 1,242 g for 5 min. The separated plasma was stored at -20°C in the laboratory until extraction. Gross morphological parameters were recorded including mass (kg), fork length (FL), and natural total length (TL), which were measured to the nearest cm over a straight line along the axis of the body from the tip of the snout to the posterior notch of the caudal fin, and to the posterior tip of the upper lobe of the caudal fin while in its natural position, respectively. Afterwards, all sharks were euthanized by severing the cervical vertebrae. Sharks captured in the Gulf of Maine, and off the coast of Rhode Island were kept in a shipboard cooler at (4°C) before transport, on ice, to the University of New England's Marine Science Center (MSC), and dissected approximately 24 h after capture. All sharks captured in the Gulf of Mexico were dissected approximately 1 h after capture. During dissections, the following were recorded: ovary and oviducal gland mass (to the nearest g); oviducal gland and the five maximum follicle diameters (to the nearest mm); and if present pup sex and stretch total length (STL), which was measured to the nearest mm over a straight line along the axis of the body from the tip of the snout to the posterior tip of the upper lobe of the caudal fin while fully extended along the axis of the body. In addition, a 5 g white skeletal muscle tissue sample from behind the second dorsal fin was collected and immediately stored at -20°C until extraction.



### **Plasma Steroid Hormone Extraction**

Each plasma sample was extracted for 17 $\beta$ -estradiol (E<sub>2</sub>), testosterone (T), and progesterone (P<sub>4</sub>) following the protocols from Tsang and Callard (1987) and Sulikowski et al. (2004). A 500  $\mu$ L aliquot of each plasma sample was extracted twice for each hormone with 10 volumes of ethyl ether (ACS grade), and the liquid phase was evaporated at 37 °C in a heat block under a stream of nitrogen. To account for procedural loss, each sample was spiked with 1,000 counts/min of tritiated [E<sub>2</sub>], [T], or [P<sub>4</sub>] (Perkin Elmer, Waltham, MA) prior to extraction. Extracts were reconstituted in phosphate buffered saline with 0.1% gelatin (PBSG).

### **Skeletal Muscle Tissue Steroid Hormone Extraction**

Two grams of white skeletal muscle tissue from each shark were homogenized with 8 mL of cold phosphate buffered saline (PBS) for 30 s using a Kinematica PT 10-35 polytron (Bohemia, NY). Then, 500  $\mu$ L aliquots of homogenate, in quadruplicate, were extracted for each hormone. Non-radiolabeled [E<sub>2</sub>], [T], and [P<sub>4</sub>] (Steraloids, Inc., Newport, RI) were used to make stocks (6.4  $\mu$ g/mL for [E<sub>2</sub>], and 80  $\mu$ g/mL for [T] and [P<sub>4</sub>]; all in ethanol, ACS grade) to “cold-spike” samples to calculate recoveries following extraction, and to generate the standard curves for radioimmunoassay. A 100  $\mu$ L aliquot of each stock was evaporated under a stream of nitrogen, reconstituted in PBSG and serially diluted to make working stocks with final concentrations of 200 pg/100 $\mu$ L for [T] and [P<sub>4</sub>], and 320 pg/100 $\mu$ L for [E<sub>2</sub>]. Working stocks were utilized to cold-spike muscle

homogenates for calculating sample recovery. Of the four homogenate replicates, two replicates were cold-spiked with a 50  $\mu$ L aliquot of the corresponding working stock (100 pg of [T] or [P<sub>4</sub>]; 160 pg of [E<sub>2</sub>]), to track sample recovery. All samples were heated in a 50 °C water bath for 15 min, briefly vortexed, and then cooled to room temperature. Samples were extracted with 10 volumes (5 mL) of 2:1 chloroform/methanol (ACS grade; Histology grade), and vortexed for 1 minute before snap freezing in a dry ice acetone bath. The liquid phase was removed by piercing the frozen homogenate layer with a pasture pipette and transferring to a new test tube. The liquid phase was then evaporated at 37 °C in a heat block under a stream of nitrogen. One mL of 70% methanol (Histology grade) was added to the dry extract, and the samples were stored at -20 °C for 24 h. Samples were centrifuged at 962 g for 10 min at 4 °C, and the liquid phase was decanted and evaporated at 37 °C in a heat block under a stream of nitrogen. Extracts were reconstituted in 200  $\mu$ L of PBSG.

Individual muscle sample recoveries were calculated by subtracting the mean concentration of hormone found in the two non-spiked replicates from the mean concentration of hormone found in the two cold-spiked replicates. The resulting concentration of hormone was then divided by the concentration of hormone that was initially added to the two cold-spiked replicates (100 pg of [T] or [P<sub>4</sub>]; 160 pg of [E<sub>2</sub>]). To calculate the final concentration of hormone in each skeletal muscle sample (in pg/g), the mean concentration of hormone found in the two non-spiked replicates was calculated, multiplied by a dilution factor of 8 (to convert units to pg/g), and finally divided by the individual sample recovery.

## **Radioimmunoassay**

Steroid hormone concentrations from both plasma and muscle extracts were determined following a modified radioimmunoassay procedure from Tsang and Callard (1987). Previously prepared non-radiolabeled [E<sub>2</sub>], [T] and [P<sub>4</sub>] stocks (6.4 µg/mL for [E<sub>2</sub>], and 80 µg/mL for [T] and [P<sub>4</sub>]) were utilized. Antibodies of [E<sub>2</sub>], [T], and [P<sub>4</sub>] (Fitzgerald Industries, Acton, MA), were diluted to final concentrations of 1:18,000, 1:10,000, and 1:2,500, respectively. Tritiated hormones, and antibodies were added to the reconstituted plasma samples (100 µL and 50 µL in duplicate) and to muscle samples (200 µL), using PBSG to bring the total volume to 400 µL. Assays were incubated at 4 °C for 24 h before separating free from bound hormone by the addition of a carbon (0.2%; Acros Organics, Fairlawn, NJ) and dextran 70 (0.02%; Tokyo Chemical Industry Co., Ltd., Tokyo, Japan) suspension followed by centrifugation at 1,242 g for 10 min at 4 °C. The resulting supernatant was then decanted into scintillation vials and 3.5 ml of Ecolume (MPO Biomedicals, Solon, OH) was added. The radioactivity level was then measured by a Perkin Elmer Tri-Carb 2900TR liquid scintillation analyzer (Waltham, MA). The mean intra-assay coefficients of variation for *S. acanthias* plasma assays were 8, 10, and 9% for [E<sub>2</sub>], [T] and [P<sub>4</sub>], respectively, and 7, 7, and 6% for *R. terraenovae* [E<sub>2</sub>], [T] and [P<sub>4</sub>] plasma assays, respectively. The mean inter-assay coefficients of variation for *S. acanthias* plasma assays were 10, 11, and 12% for [E<sub>2</sub>], [T] and [P<sub>4</sub>], respectively, and 8, 8, and 9% for *R. terraenovae* [E<sub>2</sub>], [T] and [P<sub>4</sub>] plasma assays, respectively. The mean intra-assay coefficients of variation for *S. acanthias* muscle assays were 8, 10 and 8% for [E<sub>2</sub>], [T] and [P<sub>4</sub>], respectively, and 10, 8, and 9% for *R.*

*terraenovae* [E<sub>2</sub>], [T] and [P<sub>4</sub>] muscle assays, respectively. The mean inter-assay coefficients of variation for *S. acanthias* muscle assays were 10, 11 and 12% for [E<sub>2</sub>], [T] and [P<sub>4</sub>], respectively. For calculating mean ( $\pm$  standard error) for concentrations of each steroid hormone per stage, any value that was non-detectable was assigned the lowest possible concentration that the assay would have been able to detect in the aliquots utilized.

### **Statistical Analysis**

Linear regressions were performed for [E<sub>2</sub>], [T] and [P<sub>4</sub>] quantified from plasma and skeletal muscle. Data were transformed if variables failed tests of normality or homogeneity of variance. If transformed variables still violated the assumptions the nonparametric Kendall's tau rank correlation was conducted. All data were analyzed using R 2.13.1 (R-Core Development, 2011) and graphed using means and standard error in Sigmaplot 12 (Systat Software, San Jose, CA). All tests were considered significant at  $\alpha \leq 0.05$ .

### **Results**

A total of 31 female *S. acanthias* (78-90 cm FL, 2.7-4.9 kg) were collected, and divided into discrete reproductive stages (Table 1), based on previous literature (Hisaw and Albert 1947; Tsang and Callard 1987); the group included six pre-ovulatory, six containing candles, five early gestation females containing 62-88 mm embryos, eight mid

gestation females containing 190-240 mm embryos, and six late gestation females containing 250-275 mm embryos. Ten female *R. terraenovae* (68-85 cm FL, 2.4-4.6 kg) were collected and divided into arbitrary discrete reproductive stages (Table 2), including one pre-ovulatory, two early gestation females containing 28-55 mm embryos, four early-mid gestation females containing 56-83 mm embryos, and three mid gestation females containing 85-139 mm embryos. The overall mean recoveries of plasma [E<sub>2</sub>], [T], and [P<sub>4</sub>] extractions were 80, 87, and 70% for *S. acanthias*, and 71, 85, 68% for *R. terraenovae*, respectively; while the overall mean recoveries of muscle [E<sub>2</sub>], [T], and [P<sub>4</sub>] extractions were 21, 23, and 20% for *S. acanthias*, and 59, 39, and 55% for *R. terraenovae*, respectively.

### ***S. acanthias***

In pre-ovulatory female *S. acanthias*, [P<sub>4</sub>] concentrations in both plasma (658 ± 186 pg/mL) and muscle (543 ± 285 pg/g) were similar. However, plasma [P<sub>4</sub>] (60 ± 22 pg/mL) decreased by 10-fold during the candle stage, while muscle [P<sub>4</sub>] (2412 ± 420 pg/g) increased by about 5-fold. As gestation progresses, females in early gestation displayed relatively elevated plasma (1004 ± 225 pg/mL) and muscle (1150 ± 209 pg/g) [P<sub>4</sub>] concentrations before they declined [plasma (159 ± 38 pg/mL) and muscle (386 ± 89 pg/g)] through the remainder of gestation (Figure 2a). Overall, muscle [P<sub>4</sub>] concentrations were not significantly related to plasma [P<sub>4</sub>] concentrations (Kendall's tau=0.16; p=0.23).

During the candle stage, [T] concentrations in *S. acanthias* plasma ( $23 \pm 8$  pg/mL) and muscle ( $74 \pm 24$  pg/g) were low, and they remained that way through mid gestation [plasma ( $47 \pm 11$  pg/mL) and muscle ( $134 \pm 34$  pg/g)]. However, during late gestation, plasma [T] increased ( $423 \pm 127$  pg/mL), while muscle [T] remained unchanged ( $78 \pm 18$  pg/g). The highest muscle ( $447 \pm 54$  pg/g) and plasma ( $668 \pm 255$  pg/mL) [T] concentrations were found in the pre-ovulatory stage (Figure 2b). Similar to [P<sub>4</sub>], muscle [T] was not significantly related to plasma [T] in *S. acanthias* (Kendall's tau=0.16; p=0.30).

From pre-ovulation to mid gestation, *S. acanthias* had low plasma ( $65 \pm 12$  pg/mL) and muscle ( $113 \pm 39$  pg/g) [E<sub>2</sub>] concentrations. But, concentrations of [E<sub>2</sub>] increased in plasma ( $864 \pm 116$  pg/mL) and muscle ( $364 \pm 73$  pg/g) through the second half of gestation (Figure 2c). Unlike [P<sub>4</sub>] and [T], *S. acanthias* muscle [E<sub>2</sub>] concentrations were significantly related to plasma [E<sub>2</sub>] (Kendall's tau=0.49; p=0.0017).

### **R. terraenovae**

Throughout pre-ovulation to early gestation, concentrations of [P<sub>4</sub>] and [T] in *R. terraenovae* were relatively low in plasma ([P<sub>4</sub>]:  $162 \pm 76$  pg/mL, [T]:  $151 \pm 119$  pg/mL) and muscle ([P<sub>4</sub>]:  $182 \pm 76$  pg/g, [T]:  $84 \pm 34$  pg/g). However, a marked elevation of these two hormones occurred in plasma ([P<sub>4</sub>]:  $3582 \pm 1767$  pg/mL, [T]:  $4840 \pm 884$  pg/mL) and muscle ([P<sub>4</sub>]:  $3325 \pm 1253$  pg/g, [T]:  $693 \pm 84$  pg/g) samples in early-mid gestation females. Concentrations of [P<sub>4</sub>] and [T] then decreased in the plasma ([P<sub>4</sub>]:  $132 \pm 9$  pg/mL, [T]:  $83 \pm 23$  pg/mL) and muscle ([P<sub>4</sub>]:  $391 \pm 63$  pg/g, [T]:  $141 \pm 91$  pg/g)

through mid gestation (Figure 3 a and b). Muscle [P<sub>4</sub>] and [T] of *R. terraenovae* were found to be significant predictors of plasma [P<sub>4</sub>] and [T] (linear regression: [P<sub>4</sub>] R<sup>2</sup>=0.65, p=0.0048; [T] R<sup>2</sup>=0.84, p=0.00022).

The pre-ovulatory *R. terraenovae* female contained relatively elevated concentrations of [E<sub>2</sub>] in plasma (4885 pg/mL) and muscle (232 pg/g). Concentrations of [E<sub>2</sub>] progressively decreased in both plasma (762 ± 167 pg/mL) and muscle (72 ± 12 pg/g) through early to mid gestation (Figure 3c). Similar to [P<sub>4</sub>] and [T], concentrations of *R. terraenovae* muscle [E<sub>2</sub>] were found to be significant predictors of plasma [E<sub>2</sub>] (linear regression: R<sup>2</sup>=0.79, p=0.00059).

## **Discussion**

Concentrations of reproductive hormones were successfully detected in the muscle and plasma of female *S. acanthias* and *R. terraenovae* during specific chronological events in the reproductive cycle. In addition, significant relationships were found between plasma and skeletal muscle concentrations of [E<sub>2</sub>] in *S. acanthias* and [E<sub>2</sub>], [P<sub>4</sub>], and [T] in *R. terraenovae*, suggesting that concentrations of hormones in the skeletal muscle of elasmobranchs have the potential to be an indicator of gestational stage.

### ***S. acanthias***

In the current study, plasma and muscle [P<sub>4</sub>] in *S. acanthias* exhibited similar patterns during distinct gestational stages to those previously observed by Tsang and

Callard (1987). For example, in both studies, [P<sub>4</sub>] concentrations were elevated in the first half of gestation, and began to decrease after mid gestation through parturition (Tsang and Callard 1987). Although not measured by Tsang and Callard (1987), the pre-ovulatory levels of [P<sub>4</sub>] concentrations determined in the present study were relatively elevated in both plasma and muscle when compared to the other gestational stages examined. Based on the morphological data, this increase in [P<sub>4</sub>] could inhibit further vitellogenesis (Paolucci and Callard 1998; Koob and Callard 1999), as no substantial increase in follicle diameter was noted between near parturition or pre-ovulatory females (Table 1). Follicles in this species develop concurrently with gestation, with mating and ovulation occurring shortly after birth; thus, there is presumably little follicular growth after parturition (Castro 2009). There was an observed disparity between the patterns of plasma and muscle [P<sub>4</sub>] concentrations during the candle stage of gestation. While the reasons behind this disparity cannot be determined from the current study, the differences could be attributed to [P<sub>4</sub>] uptake/clearance (D'Ercole et al. 1984), or metabolism (Payne and Hales 2004) in the plasma and skeletal muscle.

Testosterone concentrations in plasma and muscle displayed similar patterns, remaining low as gestation progressed from the candle stage through mid gestation. However, this hormone began to increase in the plasma near the end of gestation and peaked in both tissues during pre-ovulation. While there was no significant relationship between [T] concentrations in the plasma and muscle of this species, the patterns exhibited by plasma and skeletal muscle are consistent with the values observed by Tsang and Callard (1987). The functional significance of the peak in [T] during pre-ovulation has yet to be studied. It is likely not related to folliculogenesis, since follicles develop



concurrently with gestation in *S. acanthias*, with theoretically little follicular growth from parturition to pre-ovulation (Castro 2009). It is possible, however, that the peak in [T] may have a role in the storage and maintenance of sperm viability within the oviducal gland, which has been proposed in other elasmobranch species (Manire et al. 1995; Henningsen 1998).

Estradiol concentrations in the plasma and muscle of *S. acanthias* remained relatively low from pre-ovulation to mid gestation. However, substantial increases in this hormone began after mid gestation and remained elevated until just prior to parturition. These observed [E<sub>2</sub>] patterns in the current study parallel those found by Tsang and Callard (1987). However, unlike [P<sub>4</sub>] and [T] in this species, a significant relationship was found between plasma and muscle [E<sub>2</sub>] concentrations, indicating that skeletal muscle [E<sub>2</sub>] could potentially act as a reliable indicator of gestational stage in *S. acanthias*. Similar to [P<sub>4</sub>] and [T], pre-ovulatory concentrations of [E<sub>2</sub>] in *S. acanthias* have not been previously quantified. Here the present study found low concentrations in the plasma and muscle, which were expected because of the presumed minimal follicular growth after parturition (Castro 2009).

### ***R. terraenovae***

The patterns of plasma and muscle [P<sub>4</sub>] and [T] concentrations were similar in that they were relatively low from pre-ovulation through early gestation before a distinct peak was noted in early-mid gestation females. This pronounced elevation of plasma and muscle [P<sub>4</sub>] and [T] concentrations coincided with the gestational stage when uterine

compartmentalization and implantation are suggested to arise in *R. terraenovae* (Castro and Wourms 1993). Interestingly, similar findings were reported by Manire et al. (1995), who observed that [P<sub>4</sub>] and [T] concentrations were heightened in *S. tiburo* during early gestation prior to implantation, which was proposed to be a result of uterine compartmentalization (Schlernitzauer and Gilbert 1966; Callard et al. 1992). Notably, muscle [P<sub>4</sub>] and [T] concentrations were found to be significant predictors of [P<sub>4</sub>] and [T] in the plasma, suggesting that these two hormones in the skeletal muscle tissue have the potential to be indicators of gestational stage in this species.

The patterns of plasma and muscle [E<sub>2</sub>] concentrations in *R. terraenovae* also coordinately varied, being elevated during pre-ovulation before decreasing through mid gestation. This same pattern was also observed by Manire et al. (1995), who noted that in *S. tiburo*, [E<sub>2</sub>] peaks in the pre-ovulatory stage during the final phase of folliculogenesis, and decreases from early to mid gestation. In addition, muscle [E<sub>2</sub>] concentrations in *R. terraenovae* were significant predictors of plasma [E<sub>2</sub>] concentrations, indicating that skeletal muscle [E<sub>2</sub>], just like [P<sub>4</sub>] and [T], has the potential to be an indicator of gestational stage in this species.

## **Conclusions**

In the present study, we developed a method to quantify steroid hormones in skeletal muscle and tested their efficacy, when compared to plasma hormone concentrations, to assess the reproductive status of sharks. Further, an advantage of skeletal muscle over plasma is that muscle tissue can be obtained non-lethally and with

minimal invasion. The results suggest that white skeletal muscle tissue of *S. acanthias* and *R. terraenovae* is a depot for the reproductive hormones [P<sub>4</sub>], [T] and [E<sub>2</sub>], which can be reliably measured by radioimmunoassay. Additionally, the patterns of muscle reproductive hormones, coincident with gestational stage, paralleled those in plasma, despite the different reproductive modes exhibited by these two sharks. In the present study, our preliminary results thus far suggested that skeletal muscle steroid hormone content relates well to gestational stage in yolk sac placental species. As such, future *R. terraenovae* sampling efforts will be focused on collecting the later stages of gestation to determine if the parallel patterns observed in plasma and muscle steroid hormone concentrations are consistent throughout the reproductive cycle. On the other hand, while there were no significant relationships between plasma and muscle [P<sub>4</sub>] and [T] concentrations in *S. acanthias*, the similar patterns exhibited by these two hormones in plasma and skeletal muscle suggest that they could still be used to broadly assess gestational stage in *S. acanthias*. Certainly, because of the stronger relationship between plasma and skeletal muscle [E<sub>2</sub>] concentrations, this hormone may be a better indicator of gestational stage in *S. acanthias*.

### **Acknowledgements**

We thank captains C. Brown (*F/V Proud Mary*) and C. Felch (*F/V Lady Victoria*), as well as the NOAA, Pascagoula MS, bottom long line survey. Thank you Amy Carlson for creating our map. We also thank the graduate and undergraduate students of the Sulikowski lab at UNE. This project was supported by a Save our Seas Foundation grant

(reference number P170) to J.A.S, and the Marine Science Graduate Program at the University of New England. This research was conducted as part of the requirements for a master's degree at the University of New England. This manuscript represents MSC contribution number 49.

### References

- Awruch, C. A., N. W. Pankhurst, S. D. Frusher, and J. D. Stevens. 2008. Endocrine and morphological correlates of reproduction in the draughtboard shark *Cephaloscyllium laticeps* (Elasmobranchii: Scyliorhinidae). *Journal of Experimental Zoology Part A* 309A:184-197
- Barnett, L. A. K., R. L. Earley, D. A. Ebert, and G. M. Cailliet. 2009. Maturity, fecundity, and reproductive cycle of the spotted ratfish, *Hydrolagus colliei*. *Marine Biology* 156:301-316
- Callard, I. P., L. A. Fileti, L. E. Perez, L. A. Sorbera, G. Giannoukos, L. L. Klosterman, P. C. W. Tsang, and J. A. McCracken. 1992. Role of the corpus luteum and progesterone in the evolution of vertebrate viviparity. *Integrative and Comparative Biology* 32:264-275
- Castro, J. I., and J. P. Wourms. 1993. Reproduction, placentation, and embryonic development of the Atlantic sharpnose shark, *Rhizoprionodon terraenovae*. *Journal of Morphology* 218:257-280
- Castro, J. I., C. M. Woodley, and R. L. Brudek. 1999. A preliminary evaluation of the status of shark species. FAO, Technical Paper 380, Rome.

- Castro, J. I. 2009. Observations on the reproductive cycles of some viviparous North American sharks. *Aqua, International Journal of Ichthyology* 15:205-222
- D'Ercole, A. J., A. D. Stiles, and L. E. Underwood. 1984. Tissue concentrations of somatomedin C: Further evidence for multiple sites of synthesis and paracrine or autocrine mechanisms of action. *Proceedings of the National Academy of Sciences of the United States of America* 81:935-939
- Dulvy, N. K., Y. Sadovy, and J. D. Reynolds. 2003. Extinction vulnerability in marine populations. *Fish and Fisheries* 4:25-64
- Ebert, D. A. 2005. Reproductive biology of skates, *Bathyraja* (Ishiyama), along the eastern Bering Sea continental slope. *Journal of Fish Biology* 66:618-649
- Francis, M., C. O. Maolagain, and D. Stevens. 2001. Age, growth, and sexual maturity of two New Zealand endemic skates, *Dipturus nasutus* and *D. innominatus*. *New Zealand Journal of Marine and Freshwater Research* 35:831-842
- Gelsleichter, J., L. E. L. Rasmussen, C. A. Manire, J. Tyminski, B. Chang, and L. Lombardi-Carlson. 2002. Serum steroid concentrations and development of reproductive organs during puberty in male bonnethead sharks, *Sphyrna tiburo*. *Fish Physiology and Biochemistry* 26:389-401
- Hammerschlag, N., and J. Sulikowski. 2011. Killing for conservation: the need for alternatives to lethal sampling of apex predatory sharks. *Endangered Species Research* 14:135-140
- Henningsen, A. D. 1998. Levels of re-circulating reproductively-related steroid hormones in female elasmobranchs. Implications for reproduction in a captive environment. *Aquarium Sciences and Conservation* 2:97-116

- Henningsen, A. D., F. L. Murru, L. E. L. Rasmussen, B. R. Whitaker, and G. C. Violetta. 2008. Serum levels of reproductive steroid hormones in captive sand tiger sharks, *Carcharias taurus* (Rafinesque), and comments on their relation to sexual conflicts. *Fish Physiology and Biochemistry* 34:437-446
- Heupel, M. R., and C. A. Simpfendorfer. 2010. Science or slaughter: need for lethal sampling of sharks. *Conservation Biology* 24:1212-1218
- Hisaw, F. L., and A. Albert. 1947. Observations on the reproduction of the spiny dogfish, *Squalus acanthias*. *Biological Bulletin* 92:187-199
- Hoffmann, B. 1978. Use of radioimmunoassay for monitoring hormonal residues in edible animal products. *Journal Association of Official Analytical Chemists* 61:1263-1273
- Holt, W. V., and A. R. Pickard. 1999. Role of reproductive technologies and genetic resource banks in animal conservation. *Reviews of Reproduction* 4:143-150
- International Union for Conservation of Nature and Natural Resources (IUCN). 2011. IUCN Red List of Threatened Species, Version 2011.1. Available: [www.iucnredlist.org](http://www.iucnredlist.org). (July 2011).
- Kellar, N. M., M. L. Trego, C. I. Marks, and A. E. Dizon. 2006. Determining pregnancy from blubber in three species of delphinids. *Marine Mammal Science* 22:1-16
- Kneebone, J., J. A. Sulikowski, D. E. Ferguson, and P. C. W. Tsang. 2007. Endocrinological investigation into the reproductive cycles of two sympatric skate species, *Malacoraja senta* and *Amblyraja radiata*, in the western Gulf of Maine. *Environmental Biology of Fishes* 80:257-265

- Koob, T. J., and I. P. Callard. 1999. Reproductive endocrinology of female elasmobranchs: lessons from the little skate (*Raja erinacea*) and spiny dogfish (*Squalus acanthias*). *Journal of Experimental Zoology* 284:557-574
- Lasley, B. L., and J. F. Kirkpatrick. 1991. Monitoring ovarian function in captive and free-ranging wildlife by means of urinary and fecal steroids. *Journal of Zoo and Wildlife Medicine* 22:23-31
- Manire, C. A., L. E. L. Rasmussen, D. L. Hess, and H. E. Heuter. 1995. Serum steroid hormones and the reproductive cycle of the female bonnethead shark, *Sphyrna tiburo*. *General and Comparative Endocrinology* 97:366-376
- Mansour, A. A. H., D. W. McKay, J. Lien, J. C. Orr, and J. H. Banoub. 2002. Determination of pregnancy status from blubber samples in mink whales (*Balaenoptera acutorostrata*). *Marine Mammal Science* 18:112-120
- Noppe, H., B. Le Bizec, K. Verheyden, H. F. De Brabander. 2008. Novel analytical methods for the determination of steroid hormones in edible matrices. *Analytica Chimica Acta* 611:1-16
- Paolucci, M., and I. P. Callard. 1998. Characterization of progesterone-one-binding moieties in the little skate *Raja erinacea*. *General and Comparative Endocrinology* 109:106-118
- Payne, A. H., and D. B. Hales. 2004. Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocrine Reviews* 25:947-970
- Perez, L. E., and I. P. Callard. 1993. Regulation of hepatic vitellogenin synthesis in the little skate (*Raja erinacea*): Use of a homologous enzyme-linked immunosorbent assay. *Journal of Experimental Zoology* 266:31-39

- Pinhal, D., O. B. F. Gadig, A. P. Wasko, C. Oliveira, E. Ron, F. Foresti, and C. Martins. 2008. Discrimination of shark species by simple PCR of 5S rDNA repeats. *Genetics and Molecular Biology* 31:361-365
- R Development Core Team. 2011. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Available:<http://www.R-project.org/>. (September 2011).
- Ruocco, N. L., L. O. Lucifora, J. M. Díaz de Astarloa, and O. Wöhler. 2006. Reproductive biology and abundance of the white dotted skate, *Bathyraja albomaculata*, in the southwest Atlantic. *ICES Journal of Marine Science* 63:105-116
- Sawaya, W. N., K. P. Lone, A. Husain, B. Dashti, and S. Al-Zenki. 1998. Screening for estrogenic steroids in sheep and chicken by the application of enzyme-linked immunosorbent assay and a comparison with analysis by gas chromatography-mass spectrometry. *Food Chemistry* 63:563-569
- Schlernitzauer, D. A., and P. W. Gilbert. 1966. Placentation and associated aspects of gestation in the bonnethead shark, *Sphyrna tiburo*. *Journal of Morphology* 120:219-231
- Skomal, G. B. 2007. Evaluating the physiological and physical consequences of capture on post-release survivorship in large pelagic fishes. *Fisheries Management and Ecology* 14:81-89
- Sulikowski, J. A., P. C. W. Tsang, and W. H. Howell. 2004. Annual changes in steroid hormone concentrations and gonad development in the winter skate, *Leucoraja ocellata*. *Marine Biology* 144:845-853



- Sulikowski, J. A., J. Kneebone, S. Elzey, J. Jurek, W. H. Howell, and P. C. W. Tsang. 2006. Using the composite variables of reproductive morphology, histology and steroid hormones to determine age and size at sexual maturity for the thorny skate *Amblyraja radiata* in the western Gulf of Maine. *Journal of Fish Biology* 69:1449-1465
- Sulikowski, J. A., W. B. Driggers, G. W. Ingram, J. Kneebone, D. E. Ferguson, and P. C. W. Tsang. 2007a. Profiling plasma steroid hormones: a non-lethal approach for the study of skate reproductive biology and its potential use in conservation management. *Environmental Biology of Fishes* 80:285-292
- Sulikowski, J. A., S. Elzey, J. Kneebone, J. Jurek, W. H. Howell, and P. C. W. Tsang. 2007b. The reproductive cycle of the smooth skate (*Malacoraja senta*) in the western Gulf of Maine. *Journal of Marine and Freshwater Research* 58:98-103
- Tsang, P. C. W., and I. P. Callard. 1987. Morphological and endocrine correlates of the reproductive cycle of the aplacental viviparous dogfish, *Squalus acanthias*. *General and Comparative Endocrinology* 66:182-189
- Walker, T. I. 2004. Chapter 13. Management measures. Pages 285-321. *in* J. A. Musick and R. Bonfil, editors. Technical manual for the management of elasmobranchs. FAO, Fisheries Technical Paper 474, Rome.
- Walker, T. I. 2005. Reproduction in fisheries science. Pages 81-126. *in* W. C. Hamlett and B. G. M. Jamieson, editors. Reproductive biology and phylogeny of chondrichthyes (sharks, batoids and chimaeras). Science Publishers, Enfield, New Hampshire.

Walmsley-Hart, S. A., W. H. H. Sauer, and C. D. Buxton. 1999. The biology of the skates *Raja wallacei* and *R. pullopunctata* (Batoidea: Rajidae) on the Agulhas Bank, South Africa. *South African Journal of Marine Science* 21:165-179

Zeiner, S. J., and P. G. Wolf. 1993. Growth characteristics and estimates of age at maturity of two species of skates (*Raja binoculata* and *Raja rhina*) from Monterey Bay, California. Pages 87-90. *in* S. Branstetter, editor. Conservation biology of elasmobranchs. NOAA, Technical Report NMFS115, Springfield, VA.

**Table 1.** *Squalus acanthias* morphological data. Mean ( $\pm$  standard error) maximum follicle diameter (MFD) in mm, and mean ( $\pm$  standard error) pup stretch total length (STL) in mm, which was measured to the nearest mm over a straight line along the axis of the body from the tip of the snout to the posterior tip of the upper lobe of the caudal fin while fully extended along the axis of the body, by gestational stage: pre-ovulatory (P/O), candle, early, mid, and late gestation. N represents sample size by stage.

<b>Stage</b>	<b>MFD (mm)</b>	<b>Pup STL (mm)</b>	<b>N</b>
P/O	44.8 $\pm$ 1.5	N/A	5
Candle	10.0 $\pm$ 0.4	Candle	6
Early	15.6 $\pm$ 1.2	76.8 $\pm$ 5.0	5
Mid	33.8 $\pm$ 0.6	223 $\pm$ 8.3	8
Late	44.9 $\pm$ 1.2	261 $\pm$ 5.5	6

**Table 2.** *Rhizoprionodon terraenovae* morphological data. Mean ( $\pm$  standard error) maximum follicle diameter (MFD) in mm, and mean ( $\pm$  standard error) pup stretch total length (STL) in mm, which was measured to the nearest mm over a straight line along the axis of the body from the tip of the snout to the posterior tip of the upper lobe of the caudal fin while fully extended along the axis of the body, by gestational stage: pre-ovulatory (P/O), early, early-mid, and mid gestation. N represents sample size by stage.

<b>Stage</b>	<b>MFD (mm)</b>	<b>Pup STL (mm)</b>	<b>N</b>
P/O	15.6	N/A	1
Early	4.80 $\pm$ 0.9	37.7 $\pm$ 9.8	2
Early-Mid	6.15 $\pm$ 1.1	66.8 $\pm$ 4.2	4
Mid	4.37 $\pm$ 0.2	119 $\pm$ 18	3


## Figure Legends

Figure 1. Center of sampling areas of *S. acanthias*, points 1 and 2, and *R. terraenovae*, point 3.


Figure 2. Mean ( $\pm$  standard error) concentrations of plasma (pg/mL) and muscle (pg/g) steroid hormones. (A) Progesterone (B) testosterone and (C) estradiol in *Squalus acanthias*, plotted by gestational stages: pre-ovulatory (P/O), candle, early, mid, and late gestation. Numbers appearing above the x-axis represent sample size. Note the difference in scale between the plasma and muscle axes.

Figure 3. Mean ( $\pm$  standard error) concentrations of plasma (pg/mL) and muscle (pg/g) steroid hormones. (A) Progesterone (B) testosterone and (C) estradiol in *Rhizoprionodon terraenovae*, plotted by gestational stages: pre-ovulatory (P/O), early, early-mid, and mid gestation. Numbers appearing above the x-axis represent sample size. Note the difference in scale between the plasma and muscle axes.


**Figure 1**



**Figure 2**



**Figure 3**





## CHAPTER 2

### ASSESSING REPRODUCTIVE STATUS IN ELASMOBRANCH FISHES USING STEROID HORMONES EXTRACTED FROM SKELETAL MUSCLE TISSUE

#### Abstract

Elasmobranch fishes (sharks, skates, and rays) are particularly susceptible to anthropogenic threats, making a thorough understanding of their life history characteristics essential for proper management. Historically, elasmobranch reproductive data have been collected by lethal sampling, an approach that is problematic for threatened and endangered species. However, recent studies have demonstrated that non-lethal approaches can be as effective as lethal ones in assessing reproductive status. Besides plasma, we previously reported that skeletal muscle tissue, which can be obtained non-lethally and with minimal stress, is also a depot for steroid hormones. With the diverse reproductive modes exhibited by elasmobranchs, being able to determine the efficacy of skeletal muscle steroid concentrations to assess reproductive status in these fishes will enhance our knowledge of their life histories, especially for threatened and endangered species. Thus, the objective of the current study was to determine the relationship between skeletal muscle and plasma steroid hormone concentrations, along with the associated morphological changes in the reproductive tract, during the reproductive cycles in species representing the three main reproductive modes: oviparity (*Leucoraja erinacea*), aplacental viviparity (*Squalus acanthias*), and yolk-sac placental

viviparity (*Rhizoprionodon terraenovae*). The results show that concentrations of skeletal muscle progesterone (P<sub>4</sub>), testosterone (T), and estradiol (E<sub>2</sub>) were successfully quantified by radioimmunoassay in all three elasmobranch fishes. Additionally, statistically significant associations were noted between plasma and skeletal muscle P<sub>4</sub>, T and E<sub>2</sub> concentrations in *R. terraenovae* and *L. erinacea*. Furthermore, concentrations of muscle P<sub>4</sub>, T and E<sub>2</sub> were determined to be statistically significant indicators of reproductive status in the three aforementioned species. Overall, the results of the current study demonstrated that steroid hormones present in non-lethally harvested skeletal muscle tissue can be used as (strong) indicators of reproductive status and reproductive stage in elasmobranchs.

### **Introduction**

The life history characteristics of elasmobranchs, such as slow growth, and maturity late in life, make these species particularly susceptible to overexploitation (Dulvy et al., 2003). As a result, anthropogenic threats, such as direct and indirect commercial fishing, and the illegal fin trade industry, have led to significant population declines in numerous elasmobranch species (Dulvy et al., 2008). In order to manage elasmobranchs effectively, a comprehensive understanding of their life history characteristics is needed. However, data on these characteristics are lacking for many species (Castro et al., 1999; Walker, 2004; Pinhal et al., 2008; IUCN, 2011). For instance, an essential life history characteristic needed for proper management is species-specific reproductive biology, such as the onset of maturity, gestation length, and seasonal

cyclicality (Walker, 2004, 2005). When this information is incorporated into fisheries models and species assessments, insight can be gained into when, where, and how often populations are reproducing, which can ultimately aid in reducing their decline through the development of management protocols that augments fishery practices (Walker, 2004, 2005). For example, the management of blacknose sharks within the United States was directly affected and improved by the inclusion of data detailing regionally distinct reproductive periodicity (NMFS, 2011).

Lethal sampling has historically been considered the most effective approach for collecting life history information from elasmobranchs, particularly because it is quick and provides a comprehensive set of data (Heupel and Simpfendorfer, 2010). More recently, the response to species declines and ethical concerns has moved elasmobranch research towards complete preservation, which has led to the development of non-lethal sampling approaches (Sulikowski et al., 2007; Heupel and Simpfendorfer, 2010; Hammerschlag and Sulikowski, 2011).

Among the non-lethal approaches to study the reproductive biology of elasmobranchs, such as intrauterine endoscopy (Carrier et al., 2003) and ultrasonography (Daly et al., 2007), analysis of plasma hormones is the most widely used (e.g. Kneebone et al., 2007; Sulikowski et al., 2007; Henningsen et al., 2008). In particular, plasma steroid hormones have been found to correlate directly to reproductive events in elasmobranchs, such as the onset of maturity (Gelsleichter et al., 2002; Sulikowski et al., 2006) and reproductive activity (Sulikowski et al., 2004; Kneebone et al., 2007). Although plasma steroid hormone analysis has been conducted on many species of elasmobranchs spanning multiple reproductive modes (e.g. Gelsleichter et al., 2002;

Sulikowski et al., 2006; Awruch et al., 2008; Henningsen et al., 2008), obtaining blood can be stressful for the animal because of the significant amount of handling time and prolonged removal from the water (Skomal, 2007). A new non-lethal approach that is potentially less stressful and less invasive has been developed recently. Prohaska et al. (2013) investigated the efficacy of extracting steroid hormones from the skeletal muscle tissue of elasmobranchs representing two reproductive modes, the yolk sac placental viviparous *Rhizoprionodon terraenovae* and the aplacental viviparous *Squalus acanthias*. They found that steroid hormones can be successfully extracted and quantified from elasmobranch skeletal muscle tissue. But more importantly, fluctuations in these hormones may relate to gestational stage (Prohaska et al., 2013).

It is critical that effective non-lethal protocols for studying elasmobranch reproductive biology are thoroughly tested and validated prior to the shift towards their strict use in research, akin to what occurred for marine mammals in the United States (NMFS, 1972; Heupel and Simpfendorfer, 2010; Hammerschlag and Sulikowski, 2011). Because of the greater diversity of reproductive modes (yolk-sac placental viviparity, oviparity, and aplacental viviparity) in elasmobranchs and coupled with the promising results of Prohaska et al. (2013), the objectives of the present study were: 1) to attain a more comprehensive understanding of muscle steroid hormones in *R. terraenovae*; 2) to determine if this approach is appropriate for studying the reproductive biology of oviparous elasmobranchs; and 3) to determine whether skeletal muscle steroid hormones can be used as non-lethal indicators of reproductive status in elasmobranchs.

## **Methods and Materials**

### **Specimen Collection**

Female *R. terraenovae* (yolk-sac placental viviparous) and female *S. acanthias* (aplacental viviparous) were captured using the same methods, and in the same locations as previously described in Prohaska et al. (2013). Briefly, *R. terraenovae* were captured by bottom longline in the Gulf of Mexico in an area centered around 88.812 °W and 27.887 °N, while *S. acanthias* were captured by bottom trawl and gill net in the US Northwest Atlantic in an area centered around 70.115 °W and 42.471 °N. Female *L. erinacea* (oviparous) were captured by bottom trawl in November 2012 in an area centered around 70.466 °W and 42.615 °N. After capture, *L. erinacea* were maintained in an insulated 833 L live well containing ambient surface seawater. To ensure water quality was maintained, frequent water changes occurred during the approximately one and a half hour steam to the dock. All live-captured *L. erinacea* were transported from the dock in an aerated 833 L insulated live well to the University of New England's Marine Science Center (~2 h) and housed in a 3,785 L, 2.4 m diameter hexagonal tank with an open flow-through seawater system with a turnover rate of 38 L min<sup>-1</sup>. Animal husbandry of *L. erinacea* followed the protocols of Palm et al. (2011). Additionally, skates were palpated daily for up to three weeks to assess presence or absence of egg cases. Immediately prior to sampling, *L. erinacea* were euthanized by lethal pithing.

## **Sampling**

Sampling of *R. terraenovae*, *S. acanthias*, and *L. erinacea* followed the same protocols as those in Prohaska et al. (2013). Briefly, at the time of sampling, an 8 mL aliquot of blood was collected and stored at 4°C for up to 24 h. Blood was analyzed for hematocrit prior to being centrifuged at 1,242 g for 5 min. Plasma was then removed and stored at -20°C until steroid hormone analysis. The following morphological parameters were recorded: mass (kg), fork length (FL; in sharks), disc width (DW; in skates) and natural total length (TL), all of which were measured to the nearest cm over a straight line along the axis of the body. Additionally, ovary and shell gland mass (g), shell gland width (mm), and follicle diameter (mm) were recorded. From sharks, pup sex and stretch total length (STL) (mm), measured to the nearest mm over a straight line along the axis of the body, were also recorded. Most importantly, a 5 g white skeletal muscle tissue sample was collected from behind the second dorsal fin from all sharks, and from the midpoint of the right wing from all skates, and immediately stored at -20°C until analysis (Prohaska et al., 2013).

## **Plasma Steroid Hormone Extraction**

All plasma samples were extracted for progesterone (P<sub>4</sub>) testosterone (T), and 17-β-estradiol (E<sub>2</sub>) following the methods of Tsang and Callard (1987) and Sulikowski et al. (2004). Briefly, each plasma sample was extracted twice with 10 volumes of ethyl ether (ACS grade), and the organic phase was evaporated at 37 °C under a stream of nitrogen.

Extracts were reconstituted in phosphate buffered saline containing 0.1% gelatin (PBSG). Prior to extraction, each sample was spiked with 1,000 counts min<sup>-1</sup> of tritiated P<sub>4</sub>, T, or E<sub>2</sub> (Perkin Elmer, Waltham, MA) to account for procedural loss. The overall mean recoveries of P<sub>4</sub>, T, and E<sub>2</sub> were 74, 91, and 78 % for *R. terraenovae*, and 51, 83, and 60% for *L. erinacea*, respectively.

### **Skeletal Muscle Tissue Steroid Hormone Extraction**

All white skeletal muscle tissue samples were also extracted for P<sub>4</sub>, T, and E<sub>2</sub> following the protocol of Prohaska et al. (2013). Briefly, for analyzing each steroid, two grams of white skeletal muscle tissue from each individual were homogenized with 8 mL of cold phosphate buffered saline (PBS) and divided in quadruplicate. All replicates were incubated at 50 °C for 15 min, prior to extraction with 10 volumes of 2:1 chloroform/methanol (ACS grade; Histology grade). The organic phase was evaporated at 37 °C under a stream of nitrogen before reconstitution in one mL of 70% methanol (Histology grade), and kept at -20° C for 24 h. Samples were then centrifuged at 962 g for 10 min at 4 °C, before decanting and evaporating the methanol phase at 37 °C under a stream of nitrogen. Dried extracts were reconstituted in PBSG. To account for procedural loss, two of the four replicates for each muscle sample were cold-spiked with the corresponding non-radiolabelled steroid hormone (Steraloids, Inc., Newport, RI) prior to extraction. The overall mean recoveries of P<sub>4</sub>, T, and E<sub>2</sub> were 52, 47, 51% for *R. terraenovae*, and 61, 50, and 68% for *L. erinacea*, respectively.

## **Radioimmunoassay**

Plasma and muscle steroid hormone concentrations were determined by radioimmunoassay, following a modified protocol from Tsang and Callard (1987). Non-radiolabeled P<sub>4</sub>, T, and E<sub>2</sub> (Steraloids, Inc., Newport, RI) were used to make stock concentrations of 80 µg/mL for P<sub>4</sub> and T, and 6.4 µg/mL for E<sub>2</sub> in absolute ethanol (ACS grade). The P<sub>4</sub>, T and E<sub>2</sub> antibodies (Fitzgerald Industries, Acton, MA) were diluted to final concentrations of 1:2,500, 1:10,000, and 1:18,000, respectively. Tritiated hormones and antibodies were added to the reconstituted plasma and muscle samples, using PBSG to bring the total assay volume to 400 µL. After incubation at 4 °C for 24 h, free hormone was separated from bound hormone by the addition of a carbon (0.2%; Acros Organics, Fairlawn, NJ) and dextran 70 (0.02%; Tokyo Chemical Industry Co., Ltd., Tokyo, Japan) suspension, and centrifuged at 1,242 g for 10 min at 4 °C. The supernatant was combined with 3.5 mL of Ecolume (MPO Biomedicals, Solon, OH) and the radioactivity was detected by a Perkin Elmer Tri-Carb 2900TR liquid scintillation analyzer (Waltham, MA). The mean intra-assay coefficients of variation for *R. terraenovae* were 8, 7, and 5% for P<sub>4</sub>, T, and E<sub>2</sub> plasma assays, respectively, and 10, 7, 4% for *L. erinacea* P<sub>4</sub>, T, and E<sub>2</sub> plasma assays, respectively. The mean inter-assay coefficients of variation for *R. terraenovae* plasma assays were 12, 10, 12% for [P<sub>4</sub>], [T], and [E<sub>2</sub>], respectively, and 12, 12% for *L. erinacea* P<sub>4</sub>, and E<sub>2</sub> plasma assays, respectively. The mean intra-assay coefficients of variation for *R. terraenovae* were 10, 8, and 7% for P<sub>4</sub>, T, and E<sub>2</sub> muscle assays, respectively, and 5, 11, and 9% for *L. erinacea* P<sub>4</sub>, T, and E<sub>2</sub> muscle assays, respectively. The mean inter-assay coefficients of variation for *R.*



*terraenovae* were 12, 13, 12% for P<sub>4</sub>, T, and E<sub>2</sub> muscle assays, respectively, and 13, 12, 13% for *L. erinacea* P<sub>4</sub>, T, and E<sub>2</sub> muscle assays, respectively. The mean intra-assay and inter-assay coefficients of variation for *S. acanthias* can be found in Prohaska et al. (2013). When calculating mean and standard error ( $\pm$  SEM) of plasma and muscle steroid hormone concentrations per stage, any value that was non-detectable was assigned the lowest possible concentration detectable by the assay for the aliquot utilized.

### **Statistical Analysis**

Data obtained from ten *R. terraenovae* in Prohaska et al. (2013) were pooled with all *R. terraenovae* data collected in the present study for analysis. Additionally, statistical analyses were conducted on data obtained from 31 *S. acanthias* in Prohaska et al. (2013). Linear regressions were performed on plasma and skeletal muscle P<sub>4</sub>, T, and E<sub>2</sub> concentrations by species. One-way ANOVAs were performed for each species on plasma and skeletal muscle P<sub>4</sub>, T, and E<sub>2</sub> concentrations by gestational stage, followed by a Tukey's post-hoc test. If variables failed tests of normality or homogeneity of variance, the data were transformed. If transformed variables still violated the assumptions, the nonparametric Kendall's tau rank correlation or a Kruskal-Wallis rank sum test was performed instead of linear regression or one-way ANOVA, respectively. Multiple regression analyses were conducted by species to generate mathematical equations to model each of the following morphological parameters: maximum follicle diameter (MFD), ovary mass (OM), and shell gland mass (SGM) for both sharks and skates, and embryo STL for sharks, utilizing skeletal muscle P<sub>4</sub>, T, and E<sub>2</sub> concentrations as

explanatory variables. Prior to regression analyses, all skeletal muscle P<sub>4</sub>, T, and E<sub>2</sub> concentration data were log transformed to meet the assumptions of normality and homogeneity of variance. Multiple regression equations were generated by backwards selection of a fully interactive polynomial model, including all interactions between the first order hormone terms, and all first order hormone terms squared, cubed, and to the fourth power. Any term that was found to be non-significant (P>0.05) was removed from the model until all highest order terms were significant. If co-linearity was detected in a model, all first order terms within that model were centered and the multiple regression equation was re-generated following the same backwards selection procedure. All data were analyzed using R 2.15.2 (R-Core Development, 2012). All tests were considered significant at  $\alpha \leq 0.05$ .

## **Results**

### ***R. terraenovae***

A total of 24 female *R. terraenovae* (69-92 cm FL, 1.6-7.9 kg) were sampled and assigned to the following discrete reproductive stages: five immature, four pre-ovulatory, six mid gestation (containing 100-142 mm STL embryos), and nine in late gestation (containing 314-364 mm STL embryos; see Prohaska et al. 2013 and Table 1 for gestational stage details). In addition, data from 10 female *R. terraenovae* collected in Prohaska et al. (2013) were combined with the current data set, boosting the overall sample size (n=34).

Compared to mature females (Figures 1 a, b and c), immature female *R. terraenovae* had relatively low concentrations of P<sub>4</sub>, T, and E<sub>2</sub> in plasma (192 ± 46 pg/mL, n=5; 31 ± 5 pg/mL, n=5; 680 ± 140 pg/mL, n=5, respectively) and in muscle (95 ± 21 pg/g, n= 5; 50 ± 0 pg/g, n=5; 40 ± 0 pg/g, n=5, respectively).

In mature female *R. terraenovae*, from pre-ovulation to early gestation, plasma and muscle concentrations of P<sub>4</sub> and T were relatively low. During early-mid gestation, a significant increase in these hormones was observed, with plasma P<sub>4</sub> and T concentrations increasing approximately 300%, and muscle P<sub>4</sub> and T concentrations increasing approximately 800% (Figures 1 a and b). During mid gestation, P<sub>4</sub> and T concentrations significantly decreased by approximately 98% in plasma and 75% in muscle (Figures 1 a and b) (ANOVA; Plasma T: F<sub>5, 28</sub>=19.95, P=1.88e-08; Muscle T: F<sub>6, 27</sub>=6.343, P= 0.000296). During late gestation, plasma and muscle T and P<sub>4</sub> concentrations remain unchanged (Figures 1 a and b) (ANOVA; Plasma P<sub>4</sub>: F<sub>5, 28</sub>=13.64, P=8.72e-07; Muscle P<sub>4</sub>: F<sub>5, 28</sub>=14.74, P=4.12e-07). In agreement with the results of Prohaska et al. (2013), muscle P<sub>4</sub> and T were significantly correlated to plasma P<sub>4</sub> and T concentrations (P<sub>4</sub>; linear regression: R<sup>2</sup>=0.42, p=3.032e-05; T; Kendall's tau=0.43, p=0.001153).

During pre-ovulation, E<sub>2</sub> concentrations were elevated in plasma and muscle. From early to mid gestation, plasma and muscle E<sub>2</sub> concentrations significantly decreased by approximately 90%, while in late gestation, E<sub>2</sub> concentrations significantly increased by 300% in plasma and 850% in muscle (Figure 1 c) (ANOVA; Plasma E<sub>2</sub>: F<sub>5, 28</sub>=27.54, P=5.36e-10; Muscle E<sub>2</sub>: F<sub>5, 28</sub>= 62.87, P=2.38e-14). As previously reported by Prohaska

et al. (2013), *R. terraenovae* muscle E<sub>2</sub> concentrations were also found to be significantly correlated to plasma E<sub>2</sub> (linear regression: R<sup>2</sup>=0.68, p=2.041e-09).

After performing a series of multiple regression analyses on *R. terraenovae*, muscle E<sub>2</sub> concentrations are significant indicators of MFD, SGM, and OM, while a combination of muscle P<sub>4</sub> and E<sub>2</sub> concentrations are significant indicators of embryo STL (Tables 2 and 3).

### ***L. erinacea***

Fourteen female *L. erinacea* (38-59 cm TL, 25-35 cm DW, 0.34-1.09 kg) were collected and divided into the following reproductive stages: five immature, two mature non-reproductively active (MFD < 12 mm), two pre-ovulatory (MFD > 19 mm), one in the process of ovulating, with a partially formed egg case, three in pre-oviposition with fully formed egg cases, and one in post-oviposition, having laid an egg case within the previous 12 h (see Table 4 for gestational stage details) (Koob et al., 1986; Koob and Callard, 1999).

Compared to mature females (Figures 2 a, b and c), immature female *L. erinacea* had relatively low concentrations of P<sub>4</sub>, T and E<sub>2</sub> in plasma (846 ± 246 pg/mL, n=5; 710 ± 204 pg/mL, n=5; 226 ± 123 pg/mL, n=5, respectively) and in muscle (387 ± 64 pg/g, n=5; 50 ± 0 pg/g, n=5; [91 ± 18 pg/g, n=5, respectively).

Plasma and muscle P<sub>4</sub> concentrations were elevated in mature, non-reproductively active females and decreased approximately 50% into pre-ovulation (Figure 2 a). Then, P<sub>4</sub> concentrations increased by 400% and 60% in plasma and muscle, respectively, during

ovulation, before a 75% decrease in plasma and 25% decrease in muscle during pre-oviposition. Plasma and muscle P<sub>4</sub> concentrations continued to decrease by approximately 80% during post-oviposition (Figure 2 a). Similar to P<sub>4</sub> in *R. terraenovae*, *L. erinacea* muscle P<sub>4</sub> concentrations were significantly correlated to plasma P<sub>4</sub> (linear regression: R<sup>2</sup>= 0.35, p=0.02216).

In mature, non-reproductively active and pre-ovulatory females, T and E<sub>2</sub> concentrations were relatively low in plasma and muscle. An elevation in T and E<sub>2</sub> was noted during ovulation, increasing by approximately 1,500% in plasma and 500% in muscle (Figure 2 b and c). During pre-oviposition, plasma T and E<sub>2</sub> concentrations decreased by approximately 50%, while muscle T and E<sub>2</sub> increased by approximately 30%. Muscle T and E<sub>2</sub> concentrations continued to decrease by approximately 80% during post-oviposition, while plasma T increased by 66%, while plasma E<sub>2</sub> decreased by 13% (Figure 2 b and c). Similar to P<sub>4</sub>, *L. erinacea* muscle T and E<sub>2</sub> concentrations were significantly correlated to plasma T and E<sub>2</sub> (linear regression: [T] R<sup>2</sup>=0.60, p=0.001063; [E<sub>2</sub>] R<sup>2</sup>= 0.76, p=4.665e-05).

After performing multiple regression analyses on *L. erinacea*, muscle T concentrations were significant indicators of MFD, while the combination of muscle P<sub>4</sub> and E<sub>2</sub> concentrations were significant indicators of SGM, and the combination of muscle P<sub>4</sub> and T concentrations were significant indicators of OM (Tables 2 and 5).

### *S. acanthias*

A series of one-way ANOVAs were performed to investigate the relationships between P<sub>4</sub>, T, and E<sub>2</sub> and gestational stage on data previously collected from 31 female *S. acanthias* (Prohaska et al., 2013). There was a significant 350% increase in muscle P<sub>4</sub> and a significant 90% decrease in plasma P<sub>4</sub> concentrations as the sharks progressed from pre-ovulation to the candle stage of gestation (Figure 3 a). Plasma P<sub>4</sub> concentrations significantly increased by 1,500% during the early stage of gestation, while muscle P<sub>4</sub> remained relatively unchanged. From mid to late gestation, plasma and muscle P<sub>4</sub> concentrations significantly decreased by approximately 70% (ANOVA [P<sub>4</sub>] Plasma: F<sub>4, 24</sub>= 17.12, P=9.2e-07; Muscle: F<sub>4, 24</sub>=22.42, P=8.2e-08) (Figure 3 a). During pre-ovulation, muscle T concentrations were elevated, and they significantly decreased by approximately 90% in both plasma and muscle at the candle stage (Figure 3 b). Muscle T concentrations remained relatively low for the remainder of gestation, while plasma T significantly increased by 100% and 800% from mid to late gestation, respectively (ANOVA [T] Plasma: F<sub>4, 23</sub>=20.86, P=2.24e-07; Muscle: F<sub>4, 23</sub>=11.11, P=3.65e-05) (Figure 3 b). Pre-ovulatory plasma and muscle E<sub>2</sub> concentrations were relatively low, and remained unchanged until plasma concentrations significantly increased by 1,200% and muscle concentrations increased by 200% during mid-gestation (ANOVA [E<sub>2</sub>] Plasma: F<sub>4, 19</sub>=33.28, P=2.43e-08; Muscle: F<sub>4, 19</sub>=3.964, P=0.0167) (Figure 3 c).

Similar to *R. terraenovae* and *L. erinacea*, multiple regression analyses showed that a combination of muscle P<sub>4</sub>, T, and E<sub>2</sub> concentrations were significant indicators of MFD, OM, SGM, and embryo STL in *S. acanthias* (Tables 2 and 6).

## **Discussion**

In non-elasmobranch vertebrates, sex steroid hormones are successfully extracted from body depots, such as muscle, feces, and urine to study reproductive biology (e.g. Lasley and Kirkpatrick, 1991; Heppell and Sullivan, 2000; Shimizu, 2005; Barnett et al., 2009). In addition, fluctuations of the hormones present in these body depots mirror those in plasma (e.g. Hoffman and Rattenberger, 1977; Heppell and Sullivan, 2000; Shimizu, 2005; Barnett et al., 2009). For example, studies have quantified P<sub>4</sub> from blubber of marine mammals to assess if a female is mature and/or pregnant (Mansour et al., 2002; Kellar et al., 2006). Additionally, urinary and fecal E<sub>2</sub>, T, and P<sub>4</sub> have been used to track gestation in free-ranging terrestrial vertebrates for conservation purposes (Lasley and Kirkpatrick, 1991; Shimizu, 2005). In non-elasmobranch fish, fluctuations in skeletal muscle T and E<sub>2</sub> concentrations are associated with maturity, sex, and reproductive cycles (Heppell and Sullivan, 2000; Barnett et al., 2009). Thus far, previous research on a few species of elasmobranchs has reported that circulating E<sub>2</sub>, T, and P<sub>4</sub> concentrations are correlated with morphological changes in the reproductive tract and with specific events that occur during sexual maturation and the reproductive cycle (e.g. Koob et al., 1986; Rasmussen and Muru, 1992; Sulikowski et al., 2007). More recently, Prohaska et al. (2013) successfully extracted E<sub>2</sub>, T, and P<sub>4</sub> from the skeletal muscle tissue of elasmobranchs and showed that the profiles of these hormones in the muscle are similar to those in the plasma.

Steroid hormones are essential for reproduction. Plasma E<sub>2</sub> concentrations are primarily linked to the growth and maturation of follicles within the ovary of elasmobranchs (e.g. Sumpter and Dodd, 1979; Manire et al., 1995; Snelson et al., 1997, Heupel et al., 1999; Tricas et al., 2000). Medium sized follicles produce the highest concentrations of E<sub>2</sub> (Callard et al., 1993), which is transported to the liver where binding to receptors stimulates vitellogenesis (e.g. Sumpter and Dodd, 1979; Manire et al., 1995; Snelson et al., 1997, Heupel et al., 1999; Tricas et al., 2000), and subsequent accumulation of yolk within developing follicles (Perez and Callard 1989; Koob and Callard, 1999). In addition, E<sub>2</sub> also plays a prominent role in the reproductive tract (Callard et al., 1989), such as enlargement of the shell gland (Koob et al., 1986) and vascularization of the uterus (Koob and Callard, 1999). Similarly, in the current study, we found that follicular growth accompanied increases in plasma and muscle E<sub>2</sub> concentrations. For example, *R. terraenovae* plasma and muscle E<sub>2</sub> significantly increased from mid to late gestation, which corresponded to a substantial increase in follicle diameter. A similar association was observed in the yolk-sac placental viviparous species, *Sphyrna tiburo*, in which serum E<sub>2</sub> was elevated during pre-ovulation, reduced throughout the majority of gestation, and increased again during the final stages of gestation when follicular growth was beginning (Manire et al., 1995). In *S. acanthias*, an aplacental viviparous species, plasma and muscle E<sub>2</sub> concentrations increased from early to mid gestation, and remained elevated for the duration of gestation, paralleling follicular growth. These results are similar to those of Tsang and Callard (1987) who previously reported that plasma E<sub>2</sub> increased during the second half of gestation, when cohorts of follicles with larger diameters began to develop. Interestingly, we anticipated



that the highest concentration of plasma and muscle  $E_2$  in *L. erinacea* would have occurred at pre-ovulation during the period of peak follicular development (Koob et al., 1986), but this was not observed in the current study. This  $E_2$  peak might not have been detected because of the high individual variability of hormone concentrations in continuously reproducing oviparous species (Williams et al., 2013). Alternatively, the skates sampled may be at the end of their follicular growth phase and/or at the beginning of ovulatory phase.

Unlike  $E_2$ , changes in  $P_4$  are unique to reproductive mode and often linked to mode-specific reproductive events in elasmobranchs. For example, in *S. tiburo*, peaks in serum  $P_4$  concentrations during early gestation are related to uterine compartmentalization and implantation (Schlernitzauer and Gilbert, 1966; Callard et al., 1992; Manire et al., 1995). In the current study, the peak in plasma and muscle  $P_4$  during early-mid gestation *R. terraenovae* is also likely related to uterine compartmentalization and implantation, especially since implantation occurs when embryos are approximately 70-85 mm in length (Castro and Wourms, 1993). In *S. acanthias*, high concentrations of plasma  $P_4$  during early gestation are linked to embryo retention (Callard et al., 1992) and vitellogenesis suppression (Paolucci and Callard, 1998; Koob and Callard, 1999). Similar to those studies, the data herein showed that *S. acanthias* plasma and muscle  $P_4$  peaked during early gestation when little to no follicular growth was occurring, and then decreased into late gestation when follicular growth was beginning (Tsang and Callard, 1987). Despite the high individual variability in *L. erinacea* plasma  $P_4$  concentrations (Williams et al., 2013), the patterns exhibited by plasma and muscle  $P_4$  in the current study were similar to those in Koob et al. (1986). They noted that plasma  $P_4$  peaks

during ovulation, and then decreases during pre and post oviposition. In addition, Koob et al. (1986) suggested that the rise in P<sub>4</sub> might be related to ovulation and the formation of egg cases, while the decrease in P<sub>4</sub> may be related to oviposition. Further, decreases in P<sub>4</sub> after ovulation, like those found in the plasma and muscle of the current study, are suggested to inhibit the early release of egg cases by hormonally controlling and tightening cervix muscles to maintain them within the uterus to undergo sclerotization and tanning (Koob and Cox, 1988; Koob and Callard, 1999).

There is a close relationship between T and E<sub>2</sub> in aplacental viviparous and oviparous elasmobranchs. Plasma T is primarily linked to the growth of follicles (Koob et al., 1986; Tsang and Callard, 1987), with larger follicles producing the highest concentrations, which serve as a substrate for E<sub>2</sub> synthesis to facilitate vitellogenesis (Tsang and Callard, 1992) and the continued accumulation of yolk by developing follicles (Perez and Callard, 1989; Koob and Callard, 1999). In the current study, the patterns of plasma and muscle T concentrations in *S. acanthias* and *L. erinacea* were similar to those reported in the aforementioned studies, i.e. increases in T are concurrent with increases in follicle diameter (Koob et al., 1986; Tsang and Callard, 1987). However, in the yolk-sac placental viviparous *R. terraenovae*, plasma and muscle T and P<sub>4</sub> had similar profiles, with both hormones peaking during early-mid gestation. These profiles are just like the ones exhibited in *S. tiburo* serum, in which fluctuations of T were analogous to P<sub>4</sub>, peaking in early gestation when implantation and compartmentalization are suggested to occur (Manire et al., 1995).

Of the few studies so far that determined plasma sex steroid hormones in immature elasmobranchs, all have reported little to no detectable concentrations of these

hormones in fish classified as immature, based on length as well as the underdeveloped condition of external and internal reproductive organs (e.g. Rasmussen and Gruber, 1990; Rasmussen and Muru, 1992; Cicia et al., 2009). In the current study, we attempted to measure plasma and muscle P<sub>4</sub>, E<sub>2</sub>, and T concentrations in yolk-sac placental viviparous (*R. terraenovae*) and oviparous (*L. erinacea*) that were identified as immature, based on the presence of underdeveloped reproductive tracts. Similar to previous reports, concentrations of muscle and plasma P<sub>4</sub>, E<sub>2</sub>, and T in immature females were low relative to those that were mature. These results provided further support for the direct link between sex steroid hormones, the readiness of the reproductive tract and reproductive status in elasmobranchs (Rasmussen and Muru, 1992; Gelsleichter et al., 2002; Sulikowski et al., 2006; Sulikowski et al., 2007), as well as the potential use of the muscle hormones as a non-lethal method to assess maturity.

In addition to examining the relationships between muscle and plasma steroid hormones, multiple regression analyses were conducted in the current study to determine if skeletal muscle P<sub>4</sub>, T, and E<sub>2</sub> concentrations relate to SGM, OM, and MFD in *R. terraenovae*, *L. erinacea*, and *S. acanthias*, as well as to embryo STL in the two species of sharks. So far, such analyses have led to the creation of several mathematical models, although the sample sizes used to generate them were low. Nonetheless, these preliminary models are a first step towards developing the use of P<sub>4</sub>, T, and E<sub>2</sub> concentrations in non-lethally obtained skeletal muscle as indicators of reproductive status and gestational stage.

## Conclusions

The current study reported the successful detection of steroid hormones in the muscle and plasma of female *R. terraenovae*, *L. erinacea* and *S. acanthias* during specific stages of their reproductive cycles. More importantly,, plasma and skeletal muscle concentrations of P<sub>4</sub>, T, and E<sub>2</sub> in *R. terraenovae*, and *L. erinacea* were significantly associated, suggesting that skeletal muscle tissue is an appropriate substitute for plasma. The present results also affirm the efficacy of using the steroid hormones in non-lethally obtained skeletal muscle to assess reproductive status in elasmobranchs (Prohaska et al., 2013). The primary advantage of utilizing muscle versus blood is that obtaining muscle tissue may be less stressful and less invasive than obtaining blood, which will facilitate its use on large, as well as threatened species of elasmobranchs. Our previous work on *S. acanthias* and *R. terraenovae* suggests that white skeletal muscle tissue is a depot for P<sub>4</sub>, T, and E<sub>2</sub> (Prohaska et al., 2013), the present study further suggests that skeletal muscle is also a depot for the same three hormones in the oviparous *L. erinacea*. The analysis of *L. erinacea* skeletal muscle steroid hormones, so far, suggested that they may relate well to reproductive stage. However, increasing sample size will strengthen and provide a more accurate depiction of these skeletal muscle hormones throughout the reproductive cycle of an oviparous species. On the other hand, the findings of the current study provided a more complete understanding of fluctuations in P<sub>4</sub>, T, and E<sub>2</sub> during the gestation of *R. terraenovae*, reinforcing our claim that skeletal muscle steroid hormones relate well to gestation in a yolk-sac placental species.

Finally, multiple regression analyses conducted on *R. terraenovae*, *L. erinacea*, and *S. acanthias* suggest that skeletal muscle P<sub>4</sub>, T, and E<sub>2</sub> are significantly related to changes in reproductive tract morphologies during specific stages of the reproductive cycles including gestation. The robustness of these mathematical models will be strengthened by increasing sample size before they become tools for fisheries managers.

### **Acknowledgments**

We thank captains C. Brown (*F/V Proud Mary*) and C. Felch (*F/V Lady Victoria*), as well as the NOAA, Pascagoula MS, bottom longline survey. We also thank the graduate and undergraduate students of the Sulikowski lab at UNE, and the Tsang lab at UNH. This project was supported by a Save Our Seas Foundation grant (reference number P170) to J.A.S, and the Marine Science Graduate Program at the University of New England. This research was conducted as part of the requirements for a master's degree at the University of New England. This manuscript represents MSC contribution number X.

### **References**

Awruch CA, Pankhurst NW, Frusher SD, Stevens JD (2008) Endocrine and morphological correlates of reproduction in the draughtboard shark *Cephaloscyllium laticeps* (Elasmobranchii: Scyliorhinidae). *J Exp Zool Part A* 309: 184–97.

- Barnett LAK, Earley RL, Ebert DA, Cailliet GM (2009) Maturity, fecundity, and reproductive cycle of the spotted ratfish, *Hydrolagus coliei*. Mar Biol 156: 301-316.
- Callard IP, Klosterman L, Sorbera LA, Fileti LA, Reese JC (1989) Endocrine regulation of reproduction in elasmobranchs: archetype for terrestrial vertebrates. J Exp Zool Suppl 2: 12-22.
- Callard IP, Fileti LA, Perez LE, Sorbera LA, Giannoukos G, Klosterman LL, Tsang PCW, McCracken JA (1992) Role of the corpus luteum and progesterone in the evolution of vertebrate viviparity. Integr Comp Biol 32: 264-275.
- Callard IP, Koob TJ (1993) Endocrine regulation of the elasmobranch reproductive tract. J Exp Zool 266: 368-377.
- Carrier JC, Murru FL, Walsh MT, Pratt HL (2003) Assessing reproductive potential and gestation in nurse sharks (*Ginglymostoma cirratum*) using ultrasonography and endoscopy: An example of bridging the gap between field research and captive studies. Zoo Biol 22: 179-187.
- Castro JI, Woodley CM, Brudek RL (1999) A Preliminary Evaluation of the Status of Shark Species. FAO Technical paper no. 380. Rome.
- Castro JI, Wourms JP (1993) Reproduction, placentation, and embryonic development of the Atlantic sharpnose shark, *Rhizoprionodon terraenovae*. J Morphol 218: 257-280.
- Cicia AM, Driggers WB, Ingram GW, Kneebone J, Tsang PCW, Koester DM, Sulikowski JA (2009) Size and age estimates at sexual maturity for the little skate

- Leucoraja erinacea* from the western Gulf of Maine, U.S.A. J Fish Biol 75: 1648-1666.
- Daly J, Gunn I, Kirby N, Jones R, Galloway D (2007) Ultrasound examination and behavior scoring of captive broadnose sevengill sharks, *Notorynchus cepedianus*. Zoo Biol 26: 383-395.
- Dulvy N, Baum J, Clarke S, Compagno L, Cortes E, Domingo A, Fordham S, Fowler S, Francis M, Gibson C, Martinez J, Musick J, Soldo A, Stevens J, Valenti S (2008) You can swim but you can't hide: the global status and conservation of oceanic pelagic sharks and rays. Aquat Conserv 18: 459-482.
- Dulvy NK, Sadovy Y, Reynolds JD (2003) Extinction vulnerability in marine populations. Fish Fish 4: 25-64.
- Gelsleichter J, Rasmussen L, Manire C, Tyminski J, Chang B, Lombardi-Carlson L (2002) Serum steroid concentrations and development of reproductive organs during puberty in male bonnethead sharks, *Sphyrna tiburo*. Fish Physiol Biochem 26: 389-401.
- Hammerschlag N, Sulikowski J (2011) Killing for conservation: the need for alternatives to lethal sampling of apex predatory sharks. Endanger Species Res 14: 135-140.
- Henningsen AD, Murru FL, Rasmussen LEL, Whitaker BR, Violetta GC (2008) Serum levels of reproductive steroid hormones in captive sand tiger sharks, *Carcharias taurus* (Rafinesque), and comments on their relation to sexual conflicts. Fish Physiol Biochem 34: 437-46.

- Heppell S, Sullivan C (2000) Identification of gender and reproductive maturity in the absence of gonads: Muscle tissue levels of sex steroid and vitellogenin in gag (*Mycteroperca microlepis*). *Can J Fish Aquat Sci* 57: 148-159.
- Heupel MR, Whittier JM, Bennett MB (1999) Plasma steroid hormone profiles and reproductive biology of the Epaulette shark, *Hemiscyllium ocellatum*. *J Exp Zool* 284: 586-594.
- Heupel MR, Simpfendorfer CA (2010) Science or slaughter: need for lethal sampling of sharks. *Conserv Biol* 24: 1212-1218.
- Hoffmann B, Rattenberger E (1977) Testosterone concentrations in tissue from veal calves, bulls and heifers and milk-samples. *J Anim Sci*. 45: 635-641.
- International Union for Conservation of Nature and Natural Resources (IUCN). (2011) IUCN Red List of Threatened Species, Version 2011.1. <http://www.iucnredlist.org>. (last accessed July 2011).
- Kellar NM, Trego ML, Marks CI, Dizon AE (2006) Determining pregnancy from blubber in three species of delphinids. *Mar Mammal Sci* 22: 1-16.
- Kneebone J, Ferguson DE, Sulikowski JA, Tsang PCW (2007) Endocrinological investigation into the reproductive cycles of two sympatric skate species, *Malacoraja senta* and *Amblyraja radiata*, in the western Gulf of Maine. *Environ Biol Fishes* 80: 257-265.
- Koob T, Tsang PCW, Callard IP (1986) Plasma estradiol, testosterone, and progesterone levels during the ovulatory cycle of the skate (*Raja erinacea*). *Biol Reprod* 35: 267-275.



- Koob TJ, Cox DL (1988) Egg capsule catechol oxidase from the little skate *Raja erinacea* Mitchell, 1825. *Biol Bull* 175: 202-211.
- Koob TJ, Callard IP (1999) Reproductive endocrinology of female elasmobranchs: lessons from the little skate (*Raja erinacea*) and spiny dogfish (*Squalus acanthias*). *J Exp Zool* 284: 557-74.
- Lasley BL, Kirkpatrick JF (1991) Monitoring ovarian function in captive and free-ranging wildlife by means of urinary and fecal steroids. *J Zoo Wildlife Med* 22: 23-31.
- Manire CA, Rasmussen LEL, Hess DL, Heuter HE (1995) Serum steroid hormones and the reproductive cycle of the female bonnethead shark, *Sphyrna tiburo*. *Gen Comp Endocr* 97: 366-376.
- Mansour AAH, McKay DW, Lien J, Orr JC, Banoub JH (2002) Determination of pregnancy status from blubber samples in Minke whales (*Balaenoptera acutorostrata*). *Mar Mammal Sci* 18: 112-120.
- National Marine Fisheries Service (NMFS). (1972) The Marine Mammal Protection Act, as amended through 2007.
- National Marine Fisheries Service (NMFS). (2011) SEDAR 21 Stock assessment report: HMS Gulf of Mexico Blacknose Shark. US Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Office of Sustainable Fisheries, Silver Spring, MD, pp 19.
- Palm BD, Koester DM, Driggers WB, Sulikowski JA (2011) Seasonal variation in fecundity, egg case viability, gestation, and neonate size for little skates, *Leucoraja erinacea*, in the Gulf of Maine. *Environ Biol Fishes* 92: 585-589.

- Paolucci M, Callard IP (1998) Characterization of progesterone-one-binding moieties in the little skate *Raja erinacea*. Gen Comp Endocr 109: 106-118.
- Perez LE, Callard IP (1989) Evidence for progesterone inhibition of vitellogenesis in the skate, *Raja erinacea*. Amer Zool 29: 357A
- Pinhal D, Gadig OBF, Wasko AP, Oliveira C, Ron E, Foresti F, Martins C (2008) Discrimination of shark species by simple PCR of 5S rDNA repeats. Genet Mol Biol 31: 361-365.
- Prohaska BK, Tsang PCW, Driggers WB, Hoffmayer ER, Sulikowski JA (2013) Development of a non-lethal and minimally invasive protocol to study elasmobranch reproduction. Mar Coast Fish: in press.
- Rasmussen LEL, Gruber SH (1990) Serum levels of circulating steroid hormones in free-ranging carcharhinoid sharks. In: Pratt HL Jr., Gruber SH, Taniuchi T, eds, Elasmobranchs as living resources: Advances in the biology, ecology, systematics, and the status of the fisheries. NOAA technical report NMFS 90. US Department of Commerce, Springfield, IL, pp. 143-155.
- Rasmussen LEL, Murru FL (1992) Long term studies of serum concentrations of reproductively related steroid hormones in individual captive carcharhinids. Aus J Mar Fresh Res 43: 273-281.
- Schlernitzauer DA, Gilbert PW (1966) Placentation and associated aspects of gestation in the bonnethead shark, *Sphyrna tiburo*. J Morphol 120:219-231
- Shimizu K (2005) Studies on reproductive endocrinology in non-human primates: application of non-invasive methods. J Reprod Dev 51: 1-13.

- Skomal G (2007) Evaluating the physiological and physical consequences of capture on post-release survivorship in large pelagic fishes. *Fisheries Manag Ecol* 14: 81-89.
- Snelson JFF, Johnson MR, Rasmussen LEL, Hess DL (1997) Serum concentrations of steroids hormones during reproduction in the Atlantic stingray, *Dasyatis sabina*. *Gen Comp Endocr* 108: 67-79.
- Sulikowski JA, Tsang PCW, Howell HW (2004) An annual cycle of steroid hormone concentrations and gonad development in the winter skate, *Leucoraja ocellata*, from the western Gulf of Maine. *Mar Biol* 144: 845-853.
- Sulikowski JA, Kneebone J, Elzey S, Jurek J, Howell WH, Tsang PCW (2006) Using the composite variables of reproductive morphology, histology and steroid hormones to determine age and size at sexual maturity for the thorny skate *Amblyraja radiata* in the western Gulf of Maine. *J Fish Biol* 69: 1449-1465.
- Sulikowski JA, Driggers WB, Ingram GW, Kneebone J, Ferguson DE, Tsang PCW (2007) Profiling plasma steroid hormones: a non-lethal approach for the study of skate reproductive biology and its potential use in conservation management. *Environ Biol Fishes* 80: 285-292.
- Sumpter JP, Dodd JM (1979) The annual reproductive cycle of the female lesser spotted dogfish, *Scyliorhinus canicula*, and its endocrine control. *J Fish Biol* 15: 687-695.
- Tricas TC, Maruska KP, Rasmussen LEL (2000) Annual cycles of steroid hormone production, gonad development, and reproductive behavior in the Atlantic stingray. *Gen Comp Endocr* 118: 209-225.

- Tsang PCW, Callard IP (1987) Morphological and endocrine correlates of the reproductive cycle of the aplacental viviparous dogfish, *Squalus acanthias*. Gen Comp Endocr 66:182-189
- Tsang PCW, Callard IP (1992) Regulation of ovarian steroidogenesis in vitro in the viviparous shark, *Squalus acanthias*. J Exp Zool 261: 97-104.
- Walker TI (2004) Chapter 13. Management measures. In Musick JA, Bonfil R, eds, Technical manual for the management of elasmobranchs. FAO Fisheries Technical Paper no. 474. Rome, pp. 285-321.
- Walker TI (2005) Reproduction in Fisheries Science. In: Hamlett WC, Jamieson BGM, eds. Reproductive biology and phylogeny of chondrichthyes (Sharks, batoids and chimaeras). Science Publishers, Enfield, NH, pp 81-126.
- Williams LJ, Campbell MD, Tsang PCW, Sulikowski JA (2013) Using estradiol and progesterone concentrations to assess individual variability in the reproductive cyclicity of captive female little skates, *Leucoraja erinacea*, from the western Gulf of Maine. Fish Physiol Biochem. PMID, 23307173. doi:10.1007/s10695-012-9766-2

**Table 1.** *Rhizoprionodon terraenovae* morphological data. Mean ( $\pm$  standard error) of ovary mass (OM; g), shell gland mass (SGM; g), maximum follicle diameter (MFD; mm), and embryo stretch total length (STL; mm), by gestational stage: pre-ovulatory, early, early-mid, mid, and late gestation. N represents sample size.

<b>Stage</b>	<b>OM (g)</b>	<b>SGM (g)</b>	<b>MFD (mm)</b>	<b>Embryo STL (mm)</b>	<b>N</b>
Pre-ovulatory	24.3 $\pm$ 4.5	2.7 $\pm$ 0.4	20.0 $\pm$ 1.3	NA	5
Early			4.8 $\pm$ 0.9	37.7 $\pm$ 9.8	2
Early-Mid	5.8 $\pm$ 1.5	1.4 $\pm$ 0.3	6.9 $\pm$ 1.1	66.8 $\pm$ 4.2	4
Mid	3.6 $\pm$ 0.3	1.1 $\pm$ 0.2	4.3 $\pm$ 0.2	126 $\pm$ 6.8	9
Late	19.9 $\pm$ 3.2	2.9 $\pm$ 0.2	18.9 $\pm$ 1.0	338 $\pm$ 5.7	9

**Table 2.** Multiple regression analyses conducted on *Rhizoprionodon terraenovae*, *Squalus acanthias*, and *Leucoraja erinacea*. The relationships between steroid hormones and the response variables, maximum follicle diameter (MFD) (mm), embryo STL (mm), ovary mass (OM) (g), and shell gland mass (SGM) (g) are indicated by the R-squared ( $R^2$ ) and p-value. For a more detailed account of the relationships between the hormones and the response variables, see the model definitions in Tables 3, 5, 6.

	Hormones	$R^2$	p-value
<b><i>R. terraenovae</i></b>			
MFD (mm)	E <sub>2</sub>	0.900	1.18E-12
Embryo STL (mm)	E <sub>2</sub> , P <sub>4</sub>	0.843	1.52E-08
SGM (g)	E <sub>2</sub>	0.771	1.71E-08
OM (g)	E <sub>2</sub>	0.594	1.04E-05
<b><i>L. erinacea</i></b>			
SGM (g)	E <sub>2</sub> , P <sub>4</sub>	0.770	0.0115
OM (g)	P <sub>4</sub> , T	0.749	0.00869
MFD (mm)	T	0.285	0.04929
<b><i>S. acanthias</i></b>			
MFD (mm)	E <sub>2</sub> , P <sub>4</sub> , T	0.902	5.03E-06
OM (g)	E <sub>2</sub> , P <sub>4</sub> , T	0.873	0.000381
SGM (g)	E <sub>2</sub> , P <sub>4</sub> , T	0.791	1.40E-06
Embryo STL (mm)	E <sub>2</sub> , P <sub>4</sub> , T	0.747	3.81E-05

**Table 3.** Multiple regression equations generated for *Rhizoprionodon terraenovae* using skeletal muscle concentrations (pg/mL) of progesterone (P<sub>4</sub>), testosterone (T) and estradiol (E<sub>2</sub>), as indicators of the morphological characteristics: maximum follicle diameter (MFD), embryo stretch total length (STL), ovary mass (OM), and shell gland mass (SGM). The characteristics are expressed as mean (± standard error) MFD (mm), embryo STL (mm), OM (g), and SGM (g) by gestational stage. All hormone terms within the regression equations were log transformed.

Morphological Characteristic	Morphological Predictive Regression Model	Gestational Stage				
		P/O	Early	Early/Mid	Mid	Late
<b>MFD (mm)</b>	$MFD = 10.9025 + 8.0295 (E_2) + 0.7531 [(E_2)^2] - 1.1643 [(E_2)^3]^a$	20.0 ± 1.3	4.8 ± 0.9	6.9 ± 1.1	4.3 ± 0.2	18.9 ± 1.0
<b>Embryo STL (mm)</b>	$Embryo\ STL = 138.979 + 2.4.30 (E_2) - 164.588 (P_4) - 86.555 [(P_4)^2] + 35.556 [(P_4)^3] - 142.777 [(E_2)(P_4)]^a$	0	37.7 ± 9.8	66.8 ± 4.2	126.3 ± 6.8	337.7 ± 5.7
<b>OM (g)</b>	$OM = -23.398 + 6.995 (E_2)$	24.3 ± 4.5	NA	5.8 ± 1.5	3.6 ± 0.3	19.9 ± 3.2
<b>SGM (g)</b>	$SGM = -1.64748 + 0.70060 (E_2)$	2.7 ± 0.4	NA	1.4 ± 0.3	1.4 ± 0.3	2.9 ± 0.2

<sup>a</sup> All terms included in the model were centered.  
 NA- Data were not collected.

**Table 4.** *Leucoraja erinacea* morphological data. Mean ( $\pm$  standard error) of ovary mass (OM; g), maximum follicle diameter (MFD; mm), and shell gland mass (SGM; g), by reproductive stage: immature, mature (non-reproductively active), pre-ovulatory, ovulating, pre-oviposition, and post-oviposition. N represents sample size.

<b>Stage</b>	<b>OM (g)</b>	<b>MFD (mm)</b>	<b>SGM (g)</b>	<b>N</b>
Immature	1.5 $\pm$ 0.4	2.4 $\pm$ 1.0	0.5 $\pm$ 0.3	5
Mature	5.7 $\pm$ 1.2	9.5 $\pm$ 0.7	4.1 $\pm$ 0.3	2
Pre-ovulatory	5.9 $\pm$ 0.9	21 $\pm$ 1.6	5.5 $\pm$ 1.9	2
Ovulating	11.8	21.6	12.1	1
Pre-oviposition	8.6 $\pm$ 2.3	18 $\pm$ 0.9	6.7 $\pm$ 0.2	3
Post-oviposition	4.3	12.3	6.2	1



**Table 5.** Multiple regression equations generated for *L. erinacea* using skeletal muscle concentrations (pg/mL) of progesterone (P<sub>4</sub>), testosterone (T) and estradiol (E<sub>2</sub>), as indicators of morphological characteristics: maximum follicle diameter (MFD), ovary mass (OM), and shell gland mass (SGM). The characteristics are expressed as mean (± standard error) MFD (mm), OM (g), and SGM (g) by reproductive stage. All hormone terms within the regression equations were log transformed.

Morphological Characteristic	Morphological Predictive Regression Model	Reproductive Stage					
		Immature	Mature	Pre-ovulatory	Ovulating	Pre-Oviposition	Post-Oviposition
MFD (mm)	MFD = - 5.125 + 3.531 (T)	2.4 ± 1.0	9.5 ± 0.7	20.8 ± 1.6	21.6	18.3 ± 0.9	12.3
OM (g)	OM = 5.2419 + 2.1120 (E <sub>2</sub> ) + 7.8459 (P <sub>4</sub> ) + 0.6632 [(P <sub>4</sub> ) <sup>2</sup> ] - 5.9695 [(P <sub>4</sub> ) <sup>3</sup> ] <sup>a</sup>	1.5 ± 0.4	5.7 ± 1.2	5.9 ± 0.9	11.8	8.6 ± 2.3	4.3
SGM (g)	SGM = 3.7762 + 8.8563 (P <sub>4</sub> ) + 1.5244 (T) + 2.3690 [(P <sub>4</sub> ) <sup>2</sup> ] - 9.0439 [(P <sub>4</sub> ) <sup>3</sup> ] <sup>a</sup>	0.5 ± 0.3	4.1 ± 0.3	5.5 ± 1.9	12.1	6.7 ± 0.2	6.2

<sup>a</sup> All terms included in the model were centered.

**Table 6.** Multiple regression equations generated for *S. acanthias* using skeletal muscle concentrations (pg/mL) of progesterone (P<sub>4</sub>), testosterone (T) and estradiol (E<sub>2</sub>), as indicators of morphological characteristics: maximum follicle diameter (MFD), embryo stretch total length (STL), ovary mass (OM), and shell gland mass (SGM). These characteristics are expressed as mean (± standard error) hormone concentrations, MFD (mm), embryo STL (mm), OM (g), and SGM (g) by gestational stage. All hormone terms within the regression equations were log transformed and centered.

Morphological Characteristic	Morphological Predictive Regression Model	Gestational Stage				
		P/O	Candle	Early	Mid	Late
<b>MFD (mm)</b>	$MFD = -2.649 - 2.073 (E_2) + 73.115 (P_4) - 2.883 (T) - 43.726 [(P_4)^2] - 15.840 [(P_4)^3] + 10.372 [(P_4)^4] + 6.921 [(T)^2] + 3.018 [(T)(P_4)] - 5.547 [(E_2)(T)]$	44.8 ± 1.5	10.0 ± 0.4	15.6 ± 1.2	33.8 ± 0.6	44.9 ± 1.2
<b>Embryo STL (mm)</b>	$Embryo\ STL = 191.060 + 52.658 (E_2) - 58.531 (P_4) - 38.695 (T) - 28.323 [(P_4)^2] + 30.251 [(E_2)(T)]$	0	<10	76.8 ± 5.0	222.5 ± 8.1	260.7 ± 5.5
<b>OM (g)</b>	$OM = 148.648 + 1.365 (E_2) + 74.879 (P_4) - 125.265 (T) - 58.755 [(E_2)^2] + 14.208[(P_4)^2] - 25.966 [(P_4)^3] - 130.792 [(T)^2] + 105.000 [(T)^3] - 52.844 [(E_2)(T)]$	110.6 ± 9.6	28.4 ± 12.9	17.0 ± 2.3	62.9 ± 7.7	128.5 ± 21.9
<b>SGM (g)</b>	$SGM = 1.79569 - 0.26788 (E_2) - 0.32555 (P_4) + 0.15810 (T) - 0.43729 [(E_2)(T)]$	2.9 ± 0.3	1.2 ± 0.1	1.1 ± 0.1	1.3 ± 0.1	2.4 ± 0.2

## Figure Legends


Figure 1. Mean ( $\pm$  standard error) concentrations of plasma (pg/mL) and muscle (pg/g) steroid hormones. Progesterone (A), testosterone (B) and estradiol (C) in *Rhizoprionodon terraenovae*, plotted by gestational stage: immature, pre-ovulatory (P/O), early, early-mid, mid, and late gestation. Numbers above the x-axis represent sample size. Uppercase letters denote statistically significant pairwise differences in muscle hormone concentrations between gestational stages, while lowercase letters denote statistically significant pairwise differences in plasma hormone concentrations between gestational stages ( $p < 0.05$ ). Note the difference in scale between the plasma and muscle axes.

Figure 2. Mean ( $\pm$  standard error) concentrations of plasma (pg/mL) and muscle (pg/g) steroid hormones. Progesterone (A), testosterone (B) and estradiol (C) in *Leucoraja erinacea*, plotted by reproductive stage: immature, mature (non-reproductively active), pre-ovulatory, ovulating, pre-oviposition, and post-oviposition. Numbers above the x-axis represent sample size. Note the difference in scale between the plasma and muscle axes.


Figure 3. Mean ( $\pm$  standard error) concentrations of plasma (pg/mL) and muscle (pg/g) steroid hormones. Progesterone (A), testosterone (B) and estradiol (C) in *Squalus acanthias*, plotted by gestational stage: pre-ovulatory (P/O), candle, early, mid, and late gestation. Numbers above the x-axis represent sample size. Uppercase letters denote statistically significant pairwise differences in muscle hormone concentrations between

gestational stages, while lowercase letters denote statistically significant pairwise differences in plasma hormone concentrations between gestational stages ( $p < 0.05$ ). Note the difference in scale between the plasma and muscle axes.


**Figure 1.**



**Figure 2.**



**Figure 3.**



## Appendix

- Muscle extraction techniques tested. For a more detailed explanation of each technique tested please see “Bianca Prohaska Muscle Extraction Notebook #1”.
  1. 6 g of white skeletal muscle tissue homogenized with 2 mL PBS
    - a. 1 mL of homogenate extracted 3 times with 5 times the volume of ether
      - i. Reconstitute in 500 uL
        1. Low recovery
  2. 6 g of white skeletal muscle tissue homogenized with 2 mL PBS
    - a. 1 mL of homogenate extracted 3 times with 3 times the volume of ether
      - i. Reconstitute in 500 uL
        1. Low recovery
  3. 6 g of white skeletal muscle tissue homogenized with 2 mL of PBS
    - a. Centrifuge homogenate
      - i. Homogenate would not separate
  4. 3 g of white skeletal muscle tissue homogenized with 3 mL of PBS
    - a. Removed 500 uL, add 2.5 mL PBS and centrifuge
      - i. Did not separate
        1. Split remaining homogenate in 2 and dry 1 under nitrogen and 1 by air
        2. Extracted both using plasma hormone extraction technique



- a. No quantified hormone
5. 3 g of white skeletal muscle tissue homogenized with 3 mL of PBS
- a. Removed 500 uL, add 10 mL PBS and centrifuge
    - i. Did not separate
      - 1. Split remaining homogenate in 2 and dry 1 under nitrogen and 1 by air
      - 2. Extracted both using plasma hormone extraction technique
        - a. Very low concentrations of hormone
6. 2 g white skeletal muscle tissue homogenized with 8 mL of PBS
- a. 2:1 Chloroform methanol 2 rounds of extraction
    - i. Homogenate did not separate during second round of extraction, turned to slush
7. 2 g white skeletal muscle tissue homogenized with 2 mL of PBS
- a. 2:1 Chloroform methanol 1 rounds of extraction
    - i. Low recovery
8. 2 g white skeletal muscle tissue homogenized with 8 mL of PBS
- a. 2:1 Chloroform methanol 1 round of extraction
    - i. Good recoveries for estradiol and testosterone, still fairly low for progesterone, but still able to quantify hormones from all three extracts with good replication