

Gait Training a Patient who was Deaf with Multiple Total Hip Revisions: A Case Report

Natalie Slattery, BS, DPT Student; Kirsten Buchanan, PhD, PT, ATC
University of New England, Department of Physical Therapy, Portland, Maine



Background

- 93% of patients who have a total hip arthroplasty (THA) are due to end-stage osteoarthritis.
- 15-30% of patients who survive a stroke continue to live with a long-term disability.
- The most common abnormal gait pattern after surviving a stroke is due to hemiparesis.
- Patients who are deaf require greater visual and tactile cueing during gait training.
- Gait training has been shown to normalize gait patterns and increase functional mobility in patients after a total hip replacement (THR), THA revision and/or stroke.
- There is a lack of research investigating the optimal gait training plan of care (POC) for a patient with a THR, THA revision, stroke and deafness.

Purpose

The purpose of this case report was to investigate a comprehensive PT POC for a patient who is deaf and had a THR, a THA revision and a subacute stroke.

Case Description

- 77 year old woman, deaf since 8 months old
- Admitted to a skilled nursing facility after a left THA revision with residual left hemiparesis
- Required to wear an abductor brace at all times, except for skin care
- Past medical history: osteoarthritis in left and right hip and lumbar spine, elective left posterior approach THA, 4 left hip dislocations, right arterial ischemic stroke, essential HTN, stenosis of left subclavian artery
- Initial evaluation showed: decreased bilateral lower extremity strength, increase in pain level at rest and with activity, decreased balance, decreased functional mobility
- Received PT for 60-70 minutes 6-7 times/week for 4 weeks

Interventions

Gait Training	Therapeutic Exercise	Therapeutic Activity	Neuromuscular Re-Education
Amb with FWW 25 ft to 300 ft, indoor/outdoor and even/uneven surfaces	Supine: SLR, pelvic bridging	Bed Mobility: sit to and from supine & bilateral rolling	Static & dynamic balance: seated progressed to standing in // bars then to FWW
Amb without AD with hallway railing 28 ft	Seated: LAQ, hip flex, hamstring curls with 2.5 #	Transfers: sit to and from stand, SPT with FWW	Dynamic balance training during gait: stepping over and around obstacles with FWW
Ascending/descending 3 – 6 inch steps with bilateral rails	Standing: (in // bars progressed to FWW) bilateral hip abd, flex, ext, heel raises, mini squats		
Amb up/down outdoor ramp & curb step with a FWW	Standing: side stepping utilizing hallway railing, 28 ft progress to 44 ft		
Amb with FWW around fixed objects in close proximity			

Amb = ambulation, AD = assistive device, FWW = front wheeled walker, SLR = straight-leg raise, LAQ = long arc quad, # = pounds for ankle weights, // = parallel, abd = abduction, flex = flexion, ext = extension, BOS = base of support, UE = upper extremity, SPT = stand pivot transfer, ft = feet

Outcomes

Functional Mobility:	Initial Evaluation:	Discharge:
Supine to Sit	Minimal Assistance	Modified Independent
Sit to Stand	Supervision	Modified Independent
Stand Pivot Transfer	Supervision	Modified Independent
Gait	20 feet with front-wheeled walker, contact guard assistance	300 feet with front-wheeled walker, Modified Independent
Stairs	Dependent, 0 steps	Modified Independent, 3 steps
Ramp	Dependent	Supervision
CARE Item Mobility Assessment	46/84	82/84
Manual Muscle Testing	Left lower extremity 2+/5	Left lower extremity 3+/5
Numerical Pain Rating Scale	2/10 left groin at rest 6/10 left groin with activity	0/10 left groin at rest 3/10 left groin with activity

Discussion

- The patient demonstrated improvements in strength, endurance, pain level, functional mobility and gait quality.
- The patient met all short and long-term goals which allowed her to return home.
- The patient seemed to benefit from a comprehensive PT POC which focused on gait training and tactile cueing with an abductor brace.

Conclusion

- A comprehensive PT POC which focused on gait training and tactile cueing with an abductor brace was effective for a 77 year old patient who had a THR, THA revision, and a subacute stroke.
- Future research on the combined PT management of THR, THA revision, subacute stroke and deafness in a larger population of older adults is recommended.

Acknowledgments

The author acknowledges Dr. Kirsten Buchanan PhD, PT, ATC, for assistance with case report conceptualization, Jennifer Roblee PT, for supervision of the case, and the patient for their participation in the case report.

References

- Shan L, Shan B, Graham D, Saxena A. Total hip replacement: a systematic review and meta-analysis on mid-term quality of life. *Osteoarthritis Cartilage*. 2014;22:389-406. doi: 10.1016/j.joca.2013.12.006
- Xing Y, Yang SD, Dong F, et al. The beneficial role of early exercise training following stroke and possible mechanisms. *Life Sci*. 2018;198:32-37. doi: 10.1016/j.lfs.2018.02.018
- Ferrarello F, Bianchi VAM, Baccini M, et al. Tools for observational gait analysis in patients with stroke: a systematic review. *Phys Ther*. 2013;93(12):1673-1685. doi: 10.2522/ptj.20120344
- Pivec R, Johnson AJ, Mears SC, Mont MA. Hip arthroplasty. *Lancet*. 2012;380:1768-77. doi: 10.1016/S0140-6736(12)60607-2

Figure 1: Timeline of treatment




Figure 2: Patient wearing abductor brace




Figure 3: Patient walking with abductor brace using FWW