

A 6 Week Balance and Gait Training Program Using the AlterG for a Patient With Cervical Myelopathy After Spinal Decompression Surgery: A Case Report



UNIVERSITY OF
NEW ENGLAND
INNOVATION FOR A HEALTHIER PLANET

Palak Patel DPT Student

University of New England Department of Physical Therapy

Background & Purpose

- Spinal stenosis is the narrowing of the spinal canal which can cause radiculopathy or myelopathy symptoms due to compression of the spinal cord.¹
- About 80% of patients over 70 years old have some level of stenosis.¹
- There is limited research on prognosis for patients with cervical myelopathy and subsequent spinal decompression surgery.
- AlterG treadmill (AlterG Anti-Gravity Treadmill M320/F320, Fremont, CA) is an antigravity treadmill that is considered a body weight support system used in progressive gait retraining.
- The purpose of this case report is to assess the effectiveness of balance and gait training on the AlterG in a geriatric patient with severe cervical myelopathy who underwent spinal decompression surgery.

Case Description

- Patient**
 - 83-year-old male, retired farmer and lives with his wife in a ranch style home
- Aug 2015**
 - Left total hip replacement
- Nov 2016**
 - Right total hip replacement
- 2016**
 - Multiple medical complications: kidney infection, Implantable cardioverter-defibrillator, blood clots, Chronic Obstructive Pulmonary Disease and progression of decreased functional mobility
- Oct 2017**
 - Diagnosed: Cervical Stenosis with Myelopathy
 - Surgery: C4 corpectomy with C3-C5 anterior fusion and posterior cervical laminectomy and instrumented fusion
- Feb 2018**
 - The patient started Physical Therapy and Occupational Therapy services
- June 2018**
 - Patient plateaued in physical therapy
- July 2018**
 - July 2018 patient's re-evaluation
 - AlterG balance and gait training program initiated

Interventions

AlterG Ambulation: Patient ambulated in the AlterG. The Alter G has multiple settings which are adjustable to patient needs and progress.

Dual tasking: Patient ambulated in the AlterG and reached across midline for the rings and placed them on the opposite side.

Static Perturbation: Patient stands with feet together while a therapist is guarding and another therapist pulling on resistive band in various directions.

Dynamic Perturbation: Patient ambulated in the parallel bars without holding on and perturbations were given with a resistive band.

Discussion & Conclusion

- The patient demonstrated improvement in balance, gait and overall functional mobility.
- The patient improved on transfers where he only required close supervision which translated to increase independence in his own home.
- He continued physical therapy after the six-week program, to continue to address gait and balance training with the use of the AlterG.
- There is limited research on the efficacy of gait and balance training with the use of the AlterG for patients with cervical myelopathy.
- Future research should consider exploring the use of AlterG and its effects on gait and balance for geriatric patients with cervical myelopathy secondary to cervical stenosis.

Acknowledgments

The author acknowledges Matthew Somma, PT, DPT, MTC, CSCS for guidance with the conceptualization of this case report manuscript and Benjamin Kim, PT, DPT for supervision of patient care. The author acknowledges the patient for his participation in this case report.

References

1. Melancia JL, Francisco AF, Antunes JL. Spinal stenosis. In: Handbook of Clinical Neurology. Vol 119. Netherlands: Elsevier Health Sciences; 2014:541-549. doi: 10.1016/B978-0-7020-4086-3.00035-7.
2. Back & Spine Overview. Paradigm Spine. <https://www.paradigmspine.com/content/back-spine-overview>. Published March 28, 2018.

Outcomes

Outcome Measures	Week 1	Week 6
6 Minute Walk Test	Contact Guard Assist/Minimal Assistance with rolling walker 1 seated rest break Total distance: 185.62 meters	Contact Guard Assist with rolling walker 2 standing rest breaks Total distance: 264.26 meters
Berg Balance Test	28/56 (High fall risk)	35/56 (High fall risk)
30 Second Chair Stand Test	6 total stands	8 total stands